



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/57, 9/48, 1/15, 1/19, C12P 21/06, A23J 3/30	A1	(11) International Publication Number: WO 98/51803 (43) International Publication Date: 19 November 1998 (19.11.98)
(21) International Application Number: PCT/US98/09629 (22) International Filing Date: 12 May 1998 (12.05.98) (30) Priority Data: 08/857,884 16 May 1997 (16.05.97) US 60/062,892 20 October 1997 (20.10.97) US (71) Applicant: NOVO NORDISK BIOTECH, INC. [US/US]; 1445 Drew Avenue, Davis, CA 95616-4880 (US). (72) Inventors: BLINKOVSKY, Alexander; 955 Cranbrook Court #306, Davis, CA 95616 (US). BROWN, Kimberly; 8322 Windswept Court, Elk Grove, CA 95758 (US). REY, Michael, W.; 605 Robin Place, Davis, CA 95616 (US). KLOTZ, Alan; 725 Collier Drive, Dixon, CA 95620 (US). BYUN, Tony; Apartment A, 841 Adams Terrace, Davis, CA 95616 (US). (74) Agents: ZELSON, Steve, T. et al.; Novo Nordisk of North America, Inc., Suite 6400, 405 Lexington Avenue, New York, NY 10174 (US).		(81) Designated States: AL, AU, BB, BG, BR, CA, CN, CU, CZ, EE, GE, HU, IL, IS, JP, KP, KR, LC, LK, LR, LT, LU, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: POLYPEPTIDES HAVING PROLYL PIPEPTIDYL AMINOPEPTIDASE ACTIVITY AND NUCLEIC ACIDS ENCODING SAME (57) Abstract <p>The present invention relates to isolated polypeptides from <i>Aspergillus oryzae</i> having prolyl dipeptidyl aminopeptidase activity and isolated nucleic acid sequences encoding the polypeptides. The invention also relates to nucleic acid constructs, vectors, and host cells comprising the nucleic acid sequences as well as methods for producing and using the polypeptides.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

POLYPEPTIDES HAVING PROLYL PIPEPTIDYL AMINOPEPTIDASE ACTIVITY AND NUCLEIC ACIDS ENCODING SAME

5

Cross-Reference to Related Applications

This application is a continuation-in-part of pending U.S. application Serial No. 08/857,884 filed on May 16, 1997, and pending provisional U.S. application Serial No. 60/062,892 filed on October 20, 1997, which applications are fully incorporated herein by reference.

10

Background of the Invention

Field of the Invention

The present invention relates to polypeptides having dipeptidyl aminopeptidase activity and isolated nucleic acid sequences encoding the polypeptides. The invention also relates to nucleic acid constructs, vectors, and host cells comprising the nucleic acid sequences as well as methods for producing the polypeptides. The present invention further relates to methods of obtaining protein hydrolysates useful as flavour improving agents.

20

Description of the Related Art

Various food products, *e.g.*, soups, sauces and seasonings, contain flavoring agents obtained by hydrolysis of proteinaceous materials. This hydrolysis is conventionally accomplished using strong hydrochloric acid, followed by neutralization with sodium hydroxide. However, such chemical hydrolysis leads to severe degradation of the amino acids obtained during the hydrolysis, and also to hazardous byproducts formed in the course of this chemical reaction. Increasing concern over the use of flavoring agents obtained by chemical hydrolysis has led to the development of enzymatic hydrolysis processes.

25

Enzymatic hydrolysis processes of proteinaceous materials aim at obtaining a high degree of hydrolysis (DH), and this is usually attained using a complex of unspecific acting proteolytic enzymes (*i.e.*, unspecific-acting endo- and exo-peptidases). For example, WO 94/25580 describes a method for hydrolyzing proteins by use of an unspecific acting enzyme preparation obtained from *Aspergillus oryzae*. Specific acting proteolytic enzymes have not been used for this purpose because such enzymes only lead to an inadequate degree of hydrolysis.

35

Polypeptides having dipeptidyl aminopeptidase activity catalyze the removal of dipeptides from the N-terminus of peptides, polypeptides, and proteins. Such polypeptides are classified under the Enzyme Classification Number E.C. 3.4.14.- of the International Union of

Biochemistry and Molecular Biology.

Beauvais *et al.* (1997, *Journal of Biological Chemistry* 272: 6238-6244) disclose a dipeptidyl-peptidase from *Aspergillus fumigatus* which has a molecular weight of 88 kDa by SDS-PAGE and a substrate specificity limited to the hydrolysis of X-Ala, His-Ser, and Ser-Tyr dipeptides at a neutral pH optimum. Tachi *et al.* (1992, *Phytochemistry* 31: 3707-3709) disclose an X-prolyl dipeptidyl aminopeptidase from *Aspergillus oryzae* which has a molecular weight of 145 kDa by SDS-PAGE and a substrate specificity toward the peptide bond at the carboxyl site of a proline residue in the penultimate position of N-terminal free dipeptides and amides at a neutral pH optimum.

The production of protein hydrolysates with desirable organoleptic properties and high degrees of hydrolysis generally requires the use of a mixture of peptidase activities. It would be desirable to provide a single component peptidase enzyme which has activity useful for improving the organoleptic properties and degree of hydrolysis of protein hydrolysates used in food products either alone or in combination with other enzymes.

It is an object of the present invention to provide improved polypeptides having dipeptidyl aminopeptidase activity as well as methods for obtaining protein hydrolysates with desirable organoleptic qualities and high degrees of hydrolysis.

Summary of the Invention

The present invention relates to isolated polypeptides having dipeptidyl aminopeptidase activity selected from the group consisting of:

(a) a polypeptide having an amino acid sequence which has at least 50% identity with the amino acid sequence of SEQ ID NO:2;

(b) a polypeptide encoded by a nucleic acid sequence which hybridizes under medium stringency conditions with (i) the nucleic acid sequence of SEQ ID NO:1, (ii) its complementary strand, or (iii) a subsequence thereof;

(c) an allelic variant of (a) or (b); and

(d) a fragment of (a), (b), or (c), wherein the fragment has dipeptidyl aminopeptidase activity; and

(e) a polypeptide having dipeptidyl aminopeptidase activity with physicochemical properties of (i) a pH optimum in the range of from about pH 4.4 to about pH 9.8 determined after incubation for 5 minutes at ambient temperature in the presence of Ala-Pro-para-nitroanilide; (ii) a temperature stability of 90% or more, relative to initial activity, at pH 7.5 determined after incubation for 20 minutes at 65°C in the absence of substrate; and (iii) an activity towards Xaa-Pro-para-nitroanilide or Xaa-Ala-para-nitroanilide wherein Xaa is selected from the group consisting of Ala, Arg, Asp, Gly, and Val.

The present invention also relates to isolated nucleic acid sequences encoding the

polypeptides and to nucleic acid constructs, vectors, and host cells comprising the nucleic acid sequences as well as methods for producing the polypeptides.

The present invention also relates to methods for obtaining hydrolysates from proteinaceous substrates which comprise subjecting the proteinaceous material to a polypeptide
5 with dipeptidyl aminopeptidase activity alone or in combination with an endopeptidase, and to hydrolysates obtained from the method.

The present invention also relates to methods for obtaining from a proteinaceous substrate a hydrolysate enriched in free glutamic acid and/or peptide bound glutamic acid residues, which methods comprise subjecting the substrate to a deamidation process and to the
10 action of a polypeptide having dipeptidyl aminopeptidase activity.

The present invention further relates to flavor-improving compositions comprising a polypeptide with dipeptidyl aminopeptidase activity.

In a final aspect, the methods of the invention may be used in food related applications to improve flavor, such as baking. Alternatively, flavor improvement in foods may be
15 achieved by the addition of hydrolysates obtained by the methods of the invention.

Brief Description of the Figures

Figure 1 shows the nucleic acid sequence and the deduced amino acid sequence of an
20 *Aspergillus oryzae* ATCC 20386 dipeptidyl aminopeptidase (SEQ ID NOS:1 and 2, respectively).

Figure 2 shows a restriction map of pDM181.

Figure 3 shows a restriction map of pMWR54.

Detailed Description of the Invention

Polypeptides Having Dipeptidyl Aminopeptidase Activity

The term "dipeptidyl aminopeptidase activity" is defined herein as a peptidase activity which cleaves dipeptides from the N-terminal end of a peptide, polypeptide, or protein
30 sequence. Defined in a general manner, the dipeptidyl aminopeptidase is capable of cleaving the dipeptide XY from the unsubstituted N-terminal amino group of a peptide, polypeptide, or protein, wherein X or Y may represent any amino acid residue selected from the group consisting of Ala, Arg, Asn, Asp, Cys, Gln, Glu, Gly, His, Ile, Leu, Lys, Met, Phe, Pro, Ser, Thr, Trp, Tyr, and Val, but at least Ala, Arg, Asp, Gly, and/or Val. All of X and Y may
35 be different or identical. It will be understood that the isolated polypeptides having dipeptidyl aminopeptidase activity of the present invention may be unspecific as to the amino acid sequence of the dipeptide to be cleaved.

In a first embodiment, the present invention relates to isolated polypeptides having an

amino acid sequence which has a degree of identity to the amino acid sequence of SEQ ID NO:2 of at least about 50%, preferably at least about 60%, preferably at least about 70%, more preferably at least about 80%, even more preferably at least about 90%, most preferably at least about 95%, and even most preferably at least about 97%, which have dipeptidyl aminopeptidase activity (hereinafter "homologous polypeptides"). In a preferred embodiment, the homologous polypeptides have an amino acid sequence which differs by five amino acids, preferably by four amino acids, more preferably by three amino acids, even more preferably by two amino acids, and most preferably by one amino acid from the amino acid sequence of SEQ ID NO:2. For purposes of the present invention, the degree of identity between two amino acid sequences is determined by the Clustal method (Higgins, 1989, *CABIOS* 5: 151-153) with an identity table, a gap penalty of 10, and a gap length penalty of 10.

Preferably, the polypeptides of the present invention comprise the amino acid sequence of SEQ ID NO:2 or an allelic variant; and a fragment thereof, wherein the fragment has dipeptidyl aminopeptidase activity. In a more preferred embodiment, the polypeptides of the present invention comprise the amino acid sequence of SEQ ID NO:2. In another preferred embodiment, the polypeptide of the present invention has the amino acid sequence of SEQ ID NO:2 or a fragment thereof, wherein the fragment has dipeptidyl aminopeptidase activity. A fragment of SEQ ID NO:2 is a polypeptide having one or more amino acids deleted from the amino and/or carboxy terminus of this amino acid sequence. In a most preferred embodiment, the polypeptide has the amino acid sequence of SEQ ID NO:2.

Preferably, a fragment contains at least 455 amino acid residues, more preferably at least 555 amino acid residues, and most preferably at least 655 amino acid residues.

An allelic variant denotes any of two or more alternative forms of a gene occupying the same chromosomal locus. Allelic variation arises naturally through mutation, and may result in phenotypic polymorphism within populations. Gene mutations can be silent (no change in the encoded polypeptide) or may encode polypeptides having altered amino acid sequences. The term allelic variant is also used to denote a protein encoded by an allelic variant of a gene.

The amino acid sequences of the homologous polypeptides may differ from the amino acid sequence of SEQ ID NO:2 by an insertion or deletion of one or more amino acid residues and/or the substitution of one or more amino acid residues by different amino acid residues. Preferably, amino acid changes are of a minor nature, that is conservative amino acid substitutions that do not significantly affect the folding and/or activity of the protein; small deletions, typically of one to about 30 amino acids; small amino- or carboxyl-terminal extensions, such as an amino-terminal methionine residue; a small linker peptide of up to about 20-25 residues; or a small extension that facilitates purification by changing net charge or another function, such as a poly-histidine tract, an antigenic epitope or a binding domain.

Examples of conservative substitutions are within the group of basic amino acids (such as arginine, lysine and histidine), acidic amino acids (such as glutamic acid and aspartic acid),

polar amino acids (such as glutamine and asparagine), hydrophobic amino acids (such as leucine, isoleucine and valine), aromatic amino acids (such as phenylalanine, tryptophan and tyrosine), and small amino acids (such as glycine, alanine, serine, threonine and methionine). Amino acid substitutions which do not generally alter the specific activity are known in the art and are described, for example, by H. Neurath and R.L. Hill, 1979, *In, The Proteins*, Academic Press, New York. The most commonly occurring exchanges are Ala/Ser, Val/Ile, Asp/Glu, Thr/Ser, Ala/Gly, Ala/Thr, Ser/Asn, Ala/Val, Ser/Gly, Tyr/Phe, Ala/Pro, Lys/Arg, Asp/Asn, Leu/Ile, Leu/Val, Ala/Glu, and Asp/Gly as well as these in reverse.

In a second embodiment, the present invention relates to isolated polypeptides having dipeptidyl aminopeptidase activity which are encoded by nucleic acid sequences which hybridize under low stringency conditions, more preferably medium stringency conditions, and most preferably high stringency conditions, with an oligonucleotide probe which hybridizes under the same conditions with the nucleic acid sequence of SEQ ID NO:1 or its complementary strand (J. Sambrook, E.F. Fritsch, and T. Maniatus, 1989, *Molecular Cloning, A Laboratory Manual*, 2d edition, Cold Spring Harbor, New York); or allelic variants and fragments of the polypeptides, wherein the fragments have dipeptidyl aminopeptidase activity.

Hybridization indicates that the nucleic acid sequence hybridizes to the oligonucleotide probe corresponding to the polypeptide encoding part of the nucleic acid sequence shown in SEQ ID NO:1, under low to high stringency conditions (*i.e.*, prehybridization and hybridization at 42°C in 5X SSPE, 0.3% SDS, 200 µg/ml sheared and denatured salmon sperm DNA, and either 25, 35 or 50% formamide for low, medium and high stringencies, respectively), following standard Southern blotting procedures.

The amino acid sequence of SEQ ID NO:2 or a partial sequence thereof may be used to design an oligonucleotide probe, or a nucleic acid sequence encoding a polypeptide of the present invention, such as the nucleic acid sequence of SEQ ID NO:1, or a subsequence thereof, may be used to identify and clone DNA encoding polypeptides having dipeptidyl aminopeptidase activity from strains of different genera or species according to methods well known in the art. In particular, such probes can be used for hybridization with the genomic or cDNA of the genus or species of interest, following standard Southern blotting procedures, in order to identify and isolate the corresponding gene therein. Such probes can be considerably shorter than the entire sequence, but should be at least 15, preferably at least 25, and more preferably at least 40 nucleotides in length. Longer probes can also be used. Both DNA and RNA probes can be used. The probes are typically labeled for detecting the corresponding gene (for example, with ³²P, ³H, ³⁵S, biotin, or avidin).

Thus, a genomic, cDNA or combinatorial chemical library prepared from such other organisms may be screened for DNA which hybridizes with the probes described above and which encodes a polypeptide having dipeptidyl aminopeptidase activity. Genomic or other DNA from such other organisms may be separated by agarose or polyacrylamide gel

electrophoresis, or other separation techniques. DNA from the libraries or the separated DNA may be transferred to and immobilized on nitrocellulose or other suitable carrier material. In order to identify a clone or DNA which is homologous with SEQ ID NO:1, the carrier material is used in a Southern blot in which the carrier material is finally washed three times for 30 minutes each using 2 x SSC, 0.2% SDS preferably at least 50°C, more preferably at least 55°C, more preferably at least 60°C, more preferably at least 65°C, even more preferably at least 70°C, and most preferably at least 75°C. Molecules to which the oligonucleotide probe hybridizes under these conditions are detected using X-ray film.

In a third embodiment, the present invention relates to isolated polypeptides having the following physicochemical properties: (a) a pH optimum in the range of from about pH 4.4 to about pH 9.8 determined after incubation for 5 minutes at ambient temperature in the presence of Ala-Pro-para-nitroanilide; (b) a temperature stability of 90% or more, relative to initial activity, at pH 7.5 determined after incubation for 20 minutes at 65°C in the absence of substrate; and (c) an activity towards Xaa-Pro-para-nitroanilide or Xaa-Ala-para-nitroanilide wherein Xaa is selected from the group consisting of Ala, Arg, Asp, Gly, and Val. The polypeptides of the present invention also have the ability to hydrolyze other substrates.

In a preferred embodiment, the pH optimum in the range of from about pH 4.4 to about pH 9.8, more preferably in the range of from about pH 5.8 to about pH 9.8, and most preferably in the range of from about pH 7.5 to about pH 9.3 determined after incubation for 5 minutes at ambient temperature in the presence of Ala-Pro-para-nitroanilide.

In another preferred embodiment, a polypeptide of the present invention acts synergistically with an aminopeptidase to hydrolyze another polypeptide. The term "acts synergistically with an aminopeptidase to hydrolyze another polypeptide" is defined herein as the combination of a dipeptidyl aminopeptidase and an aminopeptidase which increases at least 5-fold, more preferably at least 10-fold, even more preferably at least 25-fold, most preferably at least 50-fold, and even most preferably at least 100-fold the hydrolysis of a peptide or a polypeptide relative to either individual enzyme alone. The aminopeptidase may be any aminopeptidase, but is preferably an aminopeptidase obtained from *Aspergillus oryzae*, and more preferably aminopeptidase I obtained from *Aspergillus oryzae* as described in WO 96/28542. The polypeptide is preferably a tripeptide, but may be any larger peptide or protein.

In a fourth embodiment, the present invention relates to isolated polypeptides having immunochemical identity or partial immunochemical identity to the polypeptide having the amino acid sequence of SEQ ID NO:2. The immunochemical properties are determined by immunological cross-reaction identity tests by the well-known Ouchterlony double immunodiffusion procedure. Specifically, an antiserum containing antibodies which are immunoreactive or bind to epitopes of the polypeptide having the amino acid sequence of SEQ ID NO:2 are prepared by immunizing rabbits (or other rodents) according to the procedure described by Harboe and Ingild, In N.H. Axelsen, J. Krøll, and B. Weeks, editors, *A Manual*

of *Quantitative Immunelectrophoresis*, Blackwell Scientific Publications, 1973, Chapter 23, or Johnstone and Thorpe, *Immunochemistry in Practice*, Blackwell Scientific Publications, 1982 (more specifically pages 27-31). A polypeptide having immunochemical identity is a polypeptide which reacts with the antiserum in an identical fashion such as total fusion of precipitates, identical precipitate morphology, and/or identical electrophoretic mobility using a specific immunochemical technique. A further explanation of immunochemical identity is described by Axelsen, Bock, and Krøll, In N.H. Axelsen, J. Krøll, and B. Weeks, editors, *A Manual of Quantitative Immunelectrophoresis*, Blackwell Scientific Publications, 1973, Chapter 10. A polypeptide having partial immunochemical identity is a polypeptide which reacts with the antiserum in a partially identical fashion such as partial fusion of precipitates, partially identical precipitate morphology, and/or partially identical electrophoretic mobility using a specific immunochemical technique. A further explanation of partial immunochemical identity is described by Bock and Axelsen, In N.H. Axelsen, J. Krøll, and B. Weeks, editors, *A Manual of Quantitative Immunelectrophoresis*, Blackwell Scientific Publications, 1973, Chapter 11.

Polypeptides encoded by nucleic acid sequences which hybridize with an oligonucleotide probe which hybridizes with the nucleic acid sequence of SEQ ID NO:1, its complementary strand, or allelic variants and subsequences of SEQ ID NO:1; allelic variants and fragments of the polypeptides; or the homologous polypeptides and polypeptides having identical or partially identical immunological properties may be obtained from microorganisms of any genus.

In a preferred embodiment, these polypeptides may be obtained from a bacterial source. For example, these polypeptides may be obtained from a gram positive bacterium such as a *Bacillus* strain, e.g., *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus brevis*, *Bacillus circulans*, *Bacillus coagulans*, *Bacillus lautus*, *Bacillus lentus*, *Bacillus licheniformis*, *Bacillus megaterium*, *Bacillus stearothermophilus*, *Bacillus subtilis*, or *Bacillus thuringiensis*; or a *Streptomyces* strain, e.g., *Streptomyces lividans* or *Streptomyces murinus*; or from a gram negative bacterium, e.g., *E. coli* or *Pseudomonas* sp.

The polypeptides may be obtained from a fungal source, and more preferably from a yeast strain such as a *Candida*, *Kluyveromyces*, *Pichia*, *Saccharomyces*, *Schizosaccharomyces*, or *Yarrowia* strain; or a filamentous fungal strain such as an *Acremonium*, *Aspergillus*, *Aureobasidium*, *Cryptococcus*, *Filibasidium*, *Fusarium*, *Humicola*, *Magnaporthe*, *Mucor*, *Myceliophthora*, *Neocallimastix*, *Neurospora*, *Paecilomyces*, *Penicillium*, *Piromyces*, *Schizophyllum*, *Talaromyces*, *Thermoascus*, *Thielavia*, *Tolypocladium*, or *Trichoderma* strain.

In a preferred embodiment, the polypeptides are obtained from a *Saccharomyces* *carlsbergensis*, *Saccharomyces cerevisiae*, *Saccharomyces diastaticus*, *Saccharomyces douglasii*, *Saccharomyces kluyveri*, *Saccharomyces norbensis* or *Saccharomyces oviformis*

strain.

In another preferred embodiment, the polypeptides are obtained from a *Fusarium* *bactridioides*, *Fusarium cerealis*, *Fusarium crookwellense*, *Fusarium culmorum*, *Fusarium graminearum*, *Fusarium graminum*, *Fusarium heterosporum*, *Fusarium negundi*, *Fusarium oxysporum*, *Fusarium reticulatum*, *Fusarium roseum*, *Fusarium sambucinum*, *Fusarium sarcochroum*, *Fusarium solani*, *Fusarium sporotrichioides*, *Fusarium sulphureum*, *Fusarium torulosum*, *Fusarium trichothecioides*, *Fusarium venenatum*, *Humicola insolens*, *Humicola lanuginosa*, *Mucor miehei*, *Myceliophthora thermophila*, *Neurospora crassa*, *Penicillium purpurogenum*, *Trichoderma harzianum*, *Trichoderma koningii*, *Trichoderma longibrachiatum*, *Trichoderma reesei*, or *Trichoderma viride* strain.

The polypeptides of the present invention are preferably obtained from species of *Aspergillus* including, but not limited to, *Aspergillus aculeatus*, *Aspergillus awamori*, *Aspergillus foetidus*, *Aspergillus japonicus*, *Aspergillus nidulans*, *Aspergillus niger*, or *Aspergillus oryzae*.

In a more preferred embodiment, a polypeptide of the present invention is obtained from an *Aspergillus oryzae* strain, and most preferably from *Aspergillus oryzae* ATCC 20386 or a mutant strain thereof, *e.g.*, the polypeptide with the amino acid sequence of SEQ ID NO:2.

It will be understood that for the aforementioned species, the invention encompasses both the perfect and imperfect states, and other taxonomic equivalents, *e.g.*, anamorphs, regardless of the species name by which they are known. Those skilled in the art will readily recognize the identity of appropriate equivalents. The polypeptides of the present invention may also be obtained from microorganisms which are synonyms of *Aspergillus* as defined by Raper, K.D. and Fennel, D.I., 1965, *The Genus Aspergillus*, The Wilkins Company, Baltimore. *Aspergilli* are mitosporic fungi characterized by an aspergillum comprised of a conidiospore stipe with no known teleomorphic states terminating in a vesicle, which in turn bears one or two layers of synchronously formed specialized cells, variously referred to as sterigmata or phialides, and asexually formed spores referred to as conidia. Known teleomorphs of *Aspergillus* include *Eurotium*, *Neosartorya*, and *Emericella*. Strains of *Aspergillus* and teleomorphs thereof are readily accessible to the public in a number of culture collections, such as the American Type Culture Collection (ATCC), Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (DSM), Centraalbureau Voor Schimmelcultures (CBS), and Agricultural Research Service Patent Culture Collection, Northern Regional Research Center (NRRL).

Furthermore, such polypeptides may be identified and obtained from other sources including microorganisms isolated from nature (*e.g.*, soil, composts, water, etc.) using the above-mentioned probes. Techniques for isolating microorganisms from natural habitats are well known in the art. The nucleic acid sequence may then be derived by similarly screening a genomic or cDNA library of another microorganism. Once a nucleic acid sequence encoding

the polypeptide has been detected with the probe(s), the sequence may be isolated or cloned by utilizing techniques which are known to those of ordinary skill in the art (see, *e.g.*, Sambrook *et al.*, 1989, *supra*).

For purposes of the present invention, the term "obtained from" as used herein in connection with a given source shall mean that the polypeptide is produced by the source or by a cell in which a gene from the source has been inserted.

As defined herein, an "isolated" polypeptide is a polypeptide which is essentially free of other non-dipeptidyl aminopeptidase polypeptides, *e.g.*, at least about 20% pure, preferably at least about 40% pure, more preferably about 60% pure, even more preferably about 80% pure, most preferably about 90% pure, and even most preferably about 95% pure, as determined by SDS-PAGE.

Nucleic Acid Sequences

The present invention also relates to isolated nucleic acid sequences which encode a polypeptide of the present invention. In a preferred embodiment, the nucleic acid sequence encodes a polypeptide obtained from *Aspergillus*, *e.g.*, *Aspergillus oryzae*, and in a more preferred embodiment, the nucleic acid sequence is obtained from *Aspergillus oryzae* ATCC 20386, *e.g.*, the nucleic acid sequence of SEQ ID NO:1. In another more preferred embodiment, the nucleic acid sequence is the sequence contained in plasmid pMWR52 which is contained in *Escherichia coli* NRRL B-21682. The present invention also encompasses nucleic acid sequences which encode a polypeptide having the amino acid sequence of SEQ ID NO:2, which differ from SEQ ID NO:1 by virtue of the degeneracy of the genetic code. The present invention also relates to subsequences of SEQ ID NO:1 which encode fragments of SEQ ID NO:2 which have dipeptidyl aminopeptidase activity. A subsequence of SEQ ID NO:1 is a nucleic acid sequence encompassed by SEQ ID NO:1 except that one or more nucleotides from the 5' end and/or 3' end have been deleted. Preferably, a subsequence contains at least 990 nucleotides, more preferably at least 1140 nucleotides, and most preferably at least 1290 nucleotides.

The nucleic acid sequences may be obtained from microorganisms which are taxonomic equivalents of *Aspergillus* as defined by Raper, K.D. and Fennel, D.I., 1965, *supra.*, regardless of the species name by which they are known.

The techniques used to isolate or clone a nucleic acid sequence encoding a polypeptide are known in the art and include isolation from genomic DNA, preparation from cDNA, or a combination thereof. The cloning of the nucleic acid sequences of the present invention from such genomic DNA can be effected, *e.g.*, by using the well known polymerase chain reaction (PCR) or antibody screening of expression libraries to detect cloned DNA fragments with shared structural features. See, *e.g.*, Innis *et al.*, 1990, *PCR: A Guide to Methods and Application*, Academic Press, New York. Other nucleic acid amplification procedures such as

ligase chain reaction (LCR), ligated activated transcription (LAT) and nucleic acid sequence-based amplification (NASBA) may be used. The nucleic acid sequence may be cloned from a strain of *Aspergillus*, or another or related organism and thus, for example, may be an allelic or species variant of the polypeptide encoding region of the nucleic acid sequence.

5 The term "isolated nucleic acid sequence" as used herein refers to a nucleic acid sequence which is essentially free of other nucleic acid sequences, *e.g.*, at least about 20% pure, preferably at least about 40% pure, more preferably at least about 60% pure, even more preferably at least about 80% pure, and most preferably at least about 90% pure as determined by agarose electrophoresis. For example, an isolated nucleic acid sequence can be obtained by
10 standard cloning procedures used in genetic engineering to relocate the nucleic acid sequence from its natural location to a different site where it will be reproduced. The cloning procedures may involve excision and isolation of a desired nucleic acid fragment comprising the nucleic acid sequence encoding the polypeptide, insertion of the fragment into a vector molecule, and incorporation of the recombinant vector into a host cell where multiple copies or clones of the
15 nucleic acid sequence will be replicated. The nucleic acid sequence may be of genomic, cDNA, RNA, semisynthetic, synthetic origin, or any combinations thereof.

 The present invention also relates to nucleic acid sequences which have a degree of homology to the nucleic acid sequence of SEQ ID NO:1 of at least about 50%, preferably about 60%, preferably about 70%, preferably about 80%, more preferably about 90%, even more
20 preferably about 95%, and most preferably about 97% homology, which encode an active polypeptide. For purposes of the present invention, the degree of homology between two nucleic acid sequences is determined by the Clustal method (Higgins, 1989, *supra*) with an identity table, a gap penalty of 10, and a gap length penalty of 10.

 Modification of a nucleic acid sequence encoding a polypeptide of the present invention
25 may be necessary for the synthesis of polypeptides substantially similar to the polypeptide. The term "substantially similar" to the polypeptide refers to non-naturally occurring forms of the polypeptide. These polypeptides may differ in some engineered way from the polypeptide isolated from its native source. For example, it may be of interest to synthesize variants of the polypeptide where the variants differ in specific activity, thermostability, pH optimum, or the
30 like using, *e.g.*, site-directed mutagenesis. The analogous sequence may be constructed on the basis of the nucleic acid sequence presented as the polypeptide encoding part of SEQ ID NO:1, *e.g.*, a subsequence thereof, and/or by introduction of nucleotide substitutions which do not give rise to another amino acid sequence of the polypeptide encoded by the nucleic acid sequence, but which corresponds to the codon usage of the host organism intended for
35 production of the enzyme, or by introduction of nucleotide substitutions which may give rise to a different amino acid sequence. For a general description of nucleotide substitution, see, *e.g.*, Ford *et al.*, 1991, *Protein Expression and Purification* 2: 95-107.

 It will be apparent to those skilled in the art that such substitutions can be made outside

the regions critical to the function of the molecule and still result in an active polypeptide. Amino acid residues essential to the activity of the polypeptide encoded by the isolated nucleic acid sequence of the invention, and therefore preferably not subject to substitution, may be identified according to procedures known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (see, *e.g.*, Cunningham and Wells, 1989, *Science* 244: 1081-1085). In the latter technique, mutations are introduced at every positively charged residue in the molecule, and the resultant mutant molecules are tested for dipeptidyl aminopeptidase activity to identify amino acid residues that are critical to the activity of the molecule. Sites of substrate-enzyme interaction can also be determined by analysis of the three-dimensional structure as determined by such techniques as nuclear magnetic resonance analysis, crystallography or photoaffinity labelling (see, *e.g.*, de Vos *et al.*, 1992, *Science* 255: 306-312; Smith *et al.*, 1992, *Journal of Molecular Biology* 224: 899-904; Wlodaver *et al.*, 1992, *FEBS Letters* 309: 59-64).

Polypeptides of the present invention also include fused polypeptides or cleavable fusion polypeptides in which another polypeptide is fused at the N-terminus or the C-terminus of the polypeptide or fragment thereof. A fused polypeptide is produced by fusing a nucleic acid sequence (or a portion thereof) encoding another polypeptide to a nucleic acid sequence (or a portion thereof) of the present invention. Techniques for producing fusion polypeptides are known in the art, and include ligating the coding sequences encoding the polypeptides so that they are in frame and that expression of the fused polypeptide is under control of the same promoter(s) and terminator.

The present invention also relates to isolated nucleic acid sequences encoding a polypeptide of the present invention, which hybridize under low stringency conditions, more preferably medium stringency conditions, and most preferably high stringency conditions, with an oligonucleotide probe which hybridizes under the same conditions with the nucleic acid sequence of SEQ ID NO:1 or its complementary strand; or allelic variants and subsequences thereof (Sambrook *et al.*, 1989, *supra*).

Nucleic Acid Constructs

The present invention also relates to nucleic acid constructs comprising a nucleic acid sequence of the present invention operably linked to one or more control sequences which direct the expression of the coding sequence in a suitable host cell under conditions compatible with the control sequences. Expression will be understood to include any step involved in the production of the polypeptide including, but not limited to, transcription, post-transcriptional modification, translation, post-translational modification, and secretion.

"Nucleic acid construct" is defined herein as a nucleic acid molecule, either single- or double-stranded, which is isolated from a naturally occurring gene or which has been modified to contain segments of nucleic acid which are combined and juxtaposed in a manner which

would not otherwise exist in nature. The term nucleic acid construct is synonymous with the term expression cassette when the nucleic acid construct contains all the control sequences required for expression of a coding sequence of the present invention. The term "coding sequence" as defined herein is a sequence which is transcribed into mRNA and translated into a polypeptide of the present invention. The boundaries of the coding sequence are generally determined by a ribosome binding site (prokaryotes) or by the ATG start codon (eukaryotes) located just upstream of the open reading frame at the 5' end of the mRNA and a transcription terminator sequence located just downstream of the open reading frame at the 3' end of the mRNA. A coding sequence can include, but is not limited to, DNA, cDNA, and recombinant nucleic acid sequences.

An isolated nucleic acid sequence encoding a polypeptide of the present invention may be manipulated in a variety of ways to provide for expression of the polypeptide. Manipulation of the nucleic acid sequence encoding a polypeptide prior to its insertion into a vector may be desirable or necessary depending on the expression vector. The techniques for modifying nucleic acid sequences utilizing cloning methods are well known in the art.

The term "control sequences" is defined herein to include all components which are necessary or advantageous for the expression of a polypeptide of the present invention. Each control sequence may be native or foreign to the nucleic acid sequence encoding the polypeptide. Such control sequences include, but are not limited to, a leader, a polyadenylation sequence, a propeptide sequence, a promoter, a signal sequence, and a transcription terminator.

At a minimum, the control sequences include a promoter, and transcriptional and translational stop signals. The control sequences may be provided with linkers for the purpose of introducing specific restriction sites facilitating ligation of the control sequences with the coding region of the nucleic acid sequence encoding a polypeptide. The term "operably linked" is defined herein as a configuration in which a control sequence is appropriately placed at a position relative to the coding sequence of the DNA sequence such that the control sequence directs the production of a polypeptide.

The control sequence may be an appropriate promoter sequence, a nucleic acid sequence which is recognized by a host cell for expression of the nucleic acid sequence. The promoter sequence contains transcriptional control sequences which mediate the expression of the polypeptide. The promoter may be any nucleic acid sequence which shows transcriptional activity in the host cell of choice including mutant, truncated, and hybrid promoters, and may be obtained from genes encoding extracellular or intracellular polypeptides either homologous or heterologous to the host cell.

Examples of suitable promoters for directing the transcription of the nucleic acid constructs of the present invention, especially in a bacterial host cell, are the promoters obtained from the *E. coli lac* operon, the *Streptomyces coelicolor* agarase gene (*dagA*), the *Bacillus subtilis* levansucrase gene (*sacB*), the *Bacillus licheniformis* alpha-amylase gene (*amyL*), the

Bacillus stearothermophilus maltogenic amylase gene (*amyM*), the *Bacillus amyloliquefaciens* alpha-amylase gene (*amyQ*), the *Bacillus licheniformis* penicillinase gene (*penP*), the *Bacillus subtilis* *xylA* and *xylB* genes, and the prokaryotic beta-lactamase gene (Villa-Kamaroff *et al.*, 1978, *Proceedings of the National Academy of Sciences USA* 75: 3727-3731), as well as the
5 *tac* promoter (DeBoer *et al.*, 1983, *Proceedings of the National Academy of Sciences USA* 80: 21-25). Further promoters are described in "Useful proteins from recombinant bacteria" in *Scientific American*, 1980, 242: 74-94; and in Sambrook *et al.*, 1989, *supra*.

Examples of suitable promoters for directing the transcription of the nucleic acid constructs of the present invention in a filamentous fungal host cell are promoters obtained
10 from the genes encoding *Aspergillus oryzae* TAKA amylase, *Rhizomucor miehei* aspartic proteinase, *Aspergillus niger* neutral alpha-amylase, *Aspergillus niger* acid stable alpha-amylase, *Aspergillus niger* or *Aspergillus awamori* glucoamylase (*glaA*), *Rhizomucor miehei* lipase, *Aspergillus oryzae* alkaline protease, *Aspergillus oryzae* triose phosphate isomerase, *Aspergillus nidulans* acetamidase, *Fusarium oxysporum* trypsin-like protease (U.S. Patent No.
15 4,288,627), and mutant, truncated, and hybrid promoters thereof. Particularly preferred promoters for use in filamentous fungal host cells are the TAKA amylase, NA2-tpi (a hybrid of the promoters from the genes encoding *Aspergillus niger* neutral alpha-amylase and *Aspergillus oryzae* triose phosphate isomerase), and *glaA* promoters.

In a yeast host, useful promoters are obtained from the *Saccharomyces cerevisiae*
20 enolase (ENO-1) gene, the *Saccharomyces cerevisiae* galactokinase gene (GAL1), the *Saccharomyces cerevisiae* alcohol dehydrogenase/glyceraldehyde-3-phosphate dehydrogenase genes (ADH2/GAP), and the *Saccharomyces cerevisiae* 3-phosphoglycerate kinase gene. Other useful promoters for yeast host cells are described by Romanos *et al.*, 1992, *Yeast* 8: 423-488. In a mammalian host cell, useful promoters include viral promoters such as those
25 from Simian Virus 40 (SV40), Rous sarcoma virus (RSV), adenovirus, bovine papilloma virus (BPV), and human cytomegalovirus (CMV).

The control sequence may also be a suitable transcription terminator sequence, a sequence recognized by a host cell to terminate transcription. The terminator sequence is operably linked to the 3' terminus of the nucleic acid sequence encoding the polypeptide. Any
30 terminator which is functional in the host cell of choice may be used in the present invention.

Preferred terminators for filamentous fungal host cells are obtained from the genes encoding *Aspergillus oryzae* TAKA amylase, *Aspergillus niger* glucoamylase, *Aspergillus nidulans* anthranilate synthase, *Aspergillus niger* alpha-glucosidase, and *Fusarium oxysporum* trypsin-like protease.

Preferred terminators for yeast host cells are obtained from the genes encoding *Saccharomyces cerevisiae* enolase, *Saccharomyces cerevisiae* cytochrome C (CYC1), or *Saccharomyces cerevisiae* glyceraldehyde-3-phosphate dehydrogenase. Other useful
35 terminators for yeast host cells are described by Romanos *et al.*, 1992, *supra*. Terminator

sequences are well known in the art for mammalian host cells.

The control sequence may also be a suitable leader sequence, a nontranslated region of an mRNA which is important for translation by the host cell. The leader sequence is operably linked to the 5' terminus of the nucleic acid sequence encoding the polypeptide. Any leader
5 sequence which is functional in the host cell of choice may be used in the present invention.

Preferred leaders for filamentous fungal host cells are obtained from the genes encoding *Aspergillus oryzae* TAKA amylase and *Aspergillus nidulans* triose phosphate isomerase.

Suitable leaders for yeast host cells are obtained from the *Saccharomyces cerevisiae* enolase (ENO-1) gene, the *Saccharomyces cerevisiae* 3-phosphoglycerate kinase gene, the
10 *Saccharomyces cerevisiae* alpha-factor, and the *Saccharomyces cerevisiae* alcohol dehydrogenase/glyceraldehyde-3-phosphate dehydrogenase genes (ADH2/GAP).

The control sequence may also be a polyadenylation sequence, a sequence which is operably linked to the 3' terminus of the nucleic acid sequence and which, when transcribed, is recognized by the host cell as a signal to add polyadenosine residues to transcribed mRNA.

15 Any polyadenylation sequence which is functional in the host cell of choice may be used in the present invention.

Preferred polyadenylation sequences for filamentous fungal host cells are obtained from the genes encoding *Aspergillus oryzae* TAKA amylase, *Aspergillus niger* glucoamylase, *Aspergillus nidulans* anthranilate synthase, and *Aspergillus niger* alpha-glucosidase.

20 Useful polyadenylation sequences for yeast host cells are described by Guo and Sherman, 1995, *Molecular Cellular Biology* 15: 5983-5990. Polyadenylation sequences are well known in the art for mammalian host cells.

The control sequence may also be a signal peptide coding region, which codes for an amino acid sequence linked to the amino terminus of the polypeptide which can direct the
25 encoded polypeptide into the cell's secretory pathway. The 5' end of the coding sequence of the nucleic acid sequence may inherently contain a signal peptide coding region naturally linked in translation reading frame with the segment of the coding region which encodes the secreted polypeptide. Alternatively, the 5' end of the coding sequence may contain a signal peptide coding region which is foreign to the coding sequence. The foreign signal peptide coding
30 region may be required where the coding sequence does not normally contain a signal peptide coding region. Alternatively, the foreign signal peptide coding region may simply replace the natural signal peptide coding region in order to obtain enhanced secretion of the polypeptide. The signal peptide coding region may be obtained from a glucoamylase or an amylase gene from an *Aspergillus* species, a lipase or proteinase gene from a *Rhizomucor* species, the gene
35 for the alpha-factor from *Saccharomyces cerevisiae*, an amylase or a protease gene from a *Bacillus* species, or the calf preprochymosin gene. However, any signal peptide coding region which directs the expressed polypeptide into the secretory pathway of a host cell of choice may be used in the present invention.

An effective signal peptide coding region for bacterial host cells is the signal peptide coding region obtained from the maltogenic amylase gene from *Bacillus* NCIB 11837, the *Bacillus stearothermophilus* alpha-amylase gene, the *Bacillus licheniformis* subtilisin gene, the *Bacillus licheniformis* beta-lactamase gene, the *Bacillus stearothermophilus* neutral proteases genes (*nprT*, *nprS*, *nprM*), or the *Bacillus subtilis* PrsA gene. Further signal peptides are described by Simonen and Palva, 1993, *Microbiological Reviews* 57: 109-137.

An effective signal peptide coding region for filamentous fungal host cells is the signal peptide coding region obtained from the *Aspergillus oryzae* TAKA amylase gene, *Aspergillus niger* neutral amylase gene, *Rhizomucor miehei* aspartic proteinase gene, *Humicola lanuginosa* cellulase gene, or *Humicola lanuginosa* lipase gene.

Useful signal peptides for yeast host cells are obtained from the genes for *Saccharomyces cerevisiae* alpha-factor and *Saccharomyces cerevisiae* invertase. Other useful signal peptide coding regions are described by Romanos *et al.*, 1992, *supra*.

The control sequence may also be a propeptide coding region, which codes for an amino acid sequence positioned at the amino terminus of a polypeptide. The resultant polypeptide is known as a proenzyme or propolypeptide (or a zymogen in some cases). A propolypeptide is generally inactive and can be converted to a mature active polypeptide by catalytic or autocatalytic cleavage of the propeptide from the propolypeptide. The propeptide coding region may be obtained from the *Bacillus subtilis* alkaline protease gene (*aprE*), the *Bacillus subtilis* neutral protease gene (*nprT*), the *Saccharomyces cerevisiae* alpha-factor gene, the *Rhizomucor miehei* aspartic proteinase gene, or the *Myceliophthora thermophila* laccase gene (WO 95/33836).

Where both signal peptide and propeptide regions are present at the amino terminus of a polypeptide, the propeptide region is positioned next to the amino terminus of a polypeptide and the signal peptide region is positioned next to the amino terminus of the propeptide region.

The nucleic acid constructs of the present invention may also comprise one or more nucleic acid sequences which encode one or more factors that are advantageous for directing the expression of the polypeptide, *e.g.*, a transcriptional activator (*e.g.*, a *trans*-acting factor), a chaperone, and a processing protease. Any factor that is functional in the host cell of choice may be used in the present invention. The nucleic acids encoding one or more of these factors are not necessarily in tandem with the nucleic acid sequence encoding the polypeptide.

A transcriptional activator is a protein which activates transcription of a nucleic acid sequence encoding a polypeptide (Kudla *et al.*, 1990, *EMBO Journal* 9: 1355-1364; Jarai and Buxton, 1994, *Current Genetics* 26: 2238-244; Verdier, 1990, *Yeast* 6: 271-297). The nucleic acid sequence encoding an activator may be obtained from the genes encoding *Bacillus stearothermophilus* NprA (*nprA*), *Saccharomyces cerevisiae* heme activator protein 1 (*hap1*), *Saccharomyces cerevisiae* galactose metabolizing protein 4 (*gal4*), *Aspergillus nidulans* ammonia regulation protein (*areA*), and *Aspergillus oryzae* alpha-amylase activator (*amyR*).

For further examples, see Verdier, 1990, *supra* and MacKenzie *et al.*, 1993, *Journal of General Microbiology* 139: 2295-2307.

A chaperone is a protein which assists another polypeptide in folding properly (Hartl *et al.*, 1994, *TIBS* 19: 20-25; Bergeron *et al.*, 1994, *TIBS* 19: 124-128; Demolder *et al.*, 1994, *Journal of Biotechnology* 32: 179-189; Craig, 1993, *Science* 260: 1902-1903; Gething and Sambrook, 1992, *Nature* 355: 33-45; Puig and Gilbert, 1994, *Journal of Biological Chemistry* 269: 7764-7771; Wang and Tsou, 1993, *The FASEB Journal* 7: 1515-11157; Robinson *et al.*, 1994, *Bio/Technology* 1: 381-384; Jacobs *et al.*, 1993, *Molecular Microbiology* 8: 957-966). The nucleic acid sequence encoding a chaperone may be obtained from the genes encoding *Bacillus subtilis* GroE proteins, *Bacillus subtilis* PrsA, *Aspergillus oryzae* protein disulphide isomerase, *Saccharomyces cerevisiae* calnexin, *Saccharomyces cerevisiae* BiP/GRP78, and *Saccharomyces cerevisiae* Hsp70. For further examples, see Gething and Sambrook, 1992, *supra*, and Hartl *et al.*, 1994, *supra*.

A processing protease is a protease that cleaves a propeptide to generate a mature biochemically active polypeptide (Enderlin and Ogrydziak, 1994, *Yeast* 10: 67-79; Fuller *et al.*, 1989, *Proceedings of the National Academy of Sciences USA* 86: 1434-1438; Julius *et al.*, 1984, *Cell* 37: 1075-1089; Julius *et al.*, 1983, *Cell* 32: 839-852; U.S. Patent No. 5,702,934). The nucleic acid sequence encoding a processing protease may be obtained from the genes encoding *Saccharomyces cerevisiae* dipeptidyl aminopeptidase, *Saccharomyces cerevisiae* Kex2, *Yarrowia lipolytica* dibasic processing endoprotease (*xpr6*), and *Fusarium oxysporum* metalloprotease (p45 gene).

It may also be desirable to add regulatory sequences which allow the regulation of the expression of the polypeptide relative to the growth of the host cell. Examples of regulatory systems are those which cause the expression of the gene to be turned on or off in response to a chemical or physical stimulus, including the presence of a regulatory compound. Regulatory systems in prokaryotic systems would include the *lac*, *tac*, and *trp* operator systems. In yeast, the ADH2 system or GAL1 system may be used. In filamentous fungi, the TAKA alpha-amylase promoter, *Aspergillus niger* glucoamylase promoter, and the *Aspergillus oryzae* glucoamylase promoter may be used as regulatory sequences. Other examples of regulatory sequences are those which allow for gene amplification. In eukaryotic systems, these include the dihydrofolate reductase gene which is amplified in the presence of methotrexate, and the metallothionein genes which are amplified with heavy metals. In these cases, the nucleic acid sequence encoding the polypeptide would be operably linked with the regulatory sequence.

The present invention also relates to nucleic acid constructs for altering the expression of an endogenous gene encoding a polypeptide of the present invention. The constructs may contain the minimal number of components necessary for altering expression of the endogenous gene. In one embodiment, the nucleic acid constructs preferably contain (a) a targeting sequence, (b) a regulatory sequence, (c) an exon, and (d) a splice-donor site. Upon

introduction of the nucleic acid construct into a cell, the construct inserts by homologous recombination into the cellular genome at the endogenous gene site. The targeting sequence directs the integration of elements (a)-(d) into the endogenous gene such that elements (b)-(d) are operably linked to the endogenous gene. In another embodiment, the nucleic acid constructs contain (a) a targeting sequence, (b) a regulatory sequence, (c) an exon, (d) a splice-donor site, (e) an intron, and (f) a splice-acceptor site, wherein the targeting sequence directs the integration of elements (a)-(f) such that elements (b)-(f) are operably linked to the endogenous gene. However, the constructs may contain additional components such as a selectable marker.

In both embodiments, the introduction of these components results in production of a new transcription unit in which expression of the endogenous gene is altered. In essence, the new transcription unit is a fusion product of the sequences introduced by the targeting constructs and the endogenous gene. In one embodiment in which the endogenous gene is altered, the gene is activated. In this embodiment, homologous recombination is used to replace, disrupt, or disable the regulatory region normally associated with the endogenous gene of a parent cell through the insertion of a regulatory sequence which causes the gene to be expressed at higher levels than evident in the corresponding parent cell. The activated gene can be further amplified by the inclusion of an amplifiable selectable marker gene in the construct using methods well known in the art (see, for example, U.S. Patent No. 5,641,670). In another embodiment in which the endogenous gene is altered, expression of the gene is reduced.

The targeting sequence can be within the endogenous gene, immediately adjacent to the gene, within an upstream gene, or upstream of and at a distance from the endogenous gene. One or more targeting sequences can be used. For example, a circular plasmid or DNA fragment preferably employs a single targeting sequence, while a linear plasmid or DNA fragment preferably employs two targeting sequences.

The regulatory sequence of the construct can be comprised of one or more promoters, enhancers, scaffold-attachment regions or matrix attachment sites, negative regulatory elements, transcription binding sites, or combinations of these sequences.

The constructs further contain one or more exons of the endogenous gene. An exon is defined as a DNA sequence which is copied into RNA and is present in a mature mRNA molecule such that the exon sequence is in-frame with the coding region of the endogenous gene. The exons can, optionally, contain DNA which encodes one or more amino acids and/or partially encodes an amino acid. Alternatively, the exon contains DNA which corresponds to a 5' non-encoding region. Where the exogenous exon or exons encode one or more amino acids and/or a portion of an amino acid, the nucleic acid construct is designed such that, upon transcription and splicing, the reading frame is in-frame with the coding region of the endogenous gene so that the appropriate reading frame of the portion of the mRNA derived

from the second exon is unchanged.

The splice-donor site of the constructs directs the splicing of one exon to another exon. Typically, the first exon lies 5' of the second exon, and the splice-donor site overlapping and flanking the first exon on its 3' side recognizes a splice-acceptor site flanking the second exon on the 5' side of the second exon. A splice-acceptor site, like a splice-donor site, is a sequence which directs the splicing of one exon to another exon. Acting in conjunction with a splice-donor site, the splicing apparatus uses a splice-acceptor site to effect the removal of an intron.

Expression Vectors

The present invention also relates to recombinant expression vectors comprising a nucleic acid sequence of the present invention, a promoter, and transcriptional and translational stop signals. The various nucleic acid and control sequences described above may be joined together to produce a recombinant expression vector which may include one or more convenient restriction sites to allow for insertion or substitution of the nucleic acid sequence encoding the polypeptide at such sites. Alternatively, the nucleic acid sequence of the present invention may be expressed by inserting the nucleic acid sequence or a nucleic acid construct comprising the sequence into an appropriate vector for expression. In creating the expression vector, the coding sequence is located in the vector so that the coding sequence is operably linked with the appropriate control sequences for expression, and possibly secretion.

The recombinant expression vector may be any vector (*e.g.*, a plasmid or virus) which can be conveniently subjected to recombinant DNA procedures and can bring about the expression of the nucleic acid sequence. The choice of the vector will typically depend on the compatibility of the vector with the host cell into which the vector is to be introduced. The vectors may be linear or closed circular plasmids. The vector may be an autonomously replicating vector, *i.e.*, a vector which exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, *e.g.*, a plasmid, an extrachromosomal element, a minichromosome, or an artificial chromosome. The vector may contain any means for assuring self-replication. Alternatively, the vector may be one which, when introduced into the host cell, is integrated into the genome and replicated together with the chromosome(s) into which it has been integrated. The vector system may be a single vector or plasmid or two or more vectors or plasmids which together contain the total DNA to be introduced into the genome of the host cell, or a transposon.

The vectors of the present invention preferably contain one or more selectable markers which permit easy selection of transformed cells. A selectable marker is a gene the product of which provides for biocide or viral resistance, resistance to heavy metals, prototrophy to auxotrophs, and the like. Examples of bacterial selectable markers are the *dal* genes from *Bacillus subtilis* or *Bacillus licheniformis*, or markers which confer antibiotic resistance such as ampicillin, kanamycin, chloramphenicol or tetracycline resistance. Suitable markers for

mammalian cells are the dihydrofolate reductase (*dhfr*), hygromycin phosphotransferase (*hygB*), aminoglycoside phosphotransferase II, and phleomycin resistance genes. Suitable markers for yeast host cells are ADE2, HIS3, LEU2, LYS2, MET3, TRP1, and URA3. A selectable marker for use in a filamentous fungal host cell may be selected from the group including, but not limited to, *amdS* (acetamidase), *argB* (ornithine carbamoyltransferase), *bar* (phosphinothricin acetyltransferase), *hygB* (hygromycin phosphotransferase), *niaD* (nitrate reductase), *pyrG* (orotidine-5'-phosphate decarboxylase), *sC* (sulfate adenyltransferase), *trpC* (anthranilate synthase), as well as equivalents from other species. Preferred for use in an *Aspergillus* cell are the *amdS* and *pyrG* genes of *Aspergillus nidulans* or *Aspergillus oryzae* and the *bar* gene of *Streptomyces hygroscopicus*. Furthermore, selection may be accomplished by co-transformation, *e.g.*, as described in WO 91/17243, where the selectable marker is on a separate vector.

The vectors of the present invention preferably contain an element(s) that permits stable integration of the vector into the host cell genome or autonomous replication of the vector in the cell independent of the genome of the cell.

For integration into the host cell genome, the vector may rely on the nucleic acid sequence encoding the polypeptide or any other element of the vector for stable integration of the vector into the genome by homologous or nonhomologous recombination. Alternatively, the vector may contain additional nucleic acid sequences for directing integration by homologous recombination into the genome of the host cell. The additional nucleic acid sequences enable the vector to be integrated into the host cell genome at a precise location(s) in the chromosome(s). To increase the likelihood of integration at a precise location, the integrational elements should preferably contain a sufficient number of nucleic acids, such as 100 to 1,500 base pairs, preferably 400 to 1,500 base pairs, and most preferably 800 to 1,500 base pairs, which are highly homologous with the corresponding target sequence to enhance the probability of homologous recombination. The integrational elements may be any sequence that is homologous with the target sequence in the genome of the host cell. Furthermore, the integrational elements may be non-encoding or encoding nucleic acid sequences. On the other hand, the vector may be integrated into the genome of the host cell by non-homologous recombination.

For autonomous replication, the vector may further comprise an origin of replication enabling the vector to replicate autonomously in the host cell in question. Examples of bacterial origins of replication are the origins of replication of plasmids pBR322, pUC19, pACYC177, and pACYC184 permitting replication in *E. coli*, and pUB110, pE194, pTA1060, and pAMB1 permitting replication in *Bacillus*. Examples of origins of replication for use in a yeast host cell are the 2 micron origin of replication, ARS1, ARS4, the combination of ARS1 and CEN3, and the combination of ARS4 and CEN6. The origin of replication may be one having a mutation which makes its functioning temperature-sensitive in the host cell (see, *e.g.*, Ehrlich, 1978,

Proceedings of the National Academy of Sciences USA 75: 1433).

More than one copy of a nucleic acid sequence encoding a polypeptide of the present invention may be inserted into the host cell to amplify expression of the nucleic acid sequence. Stable amplification of the nucleic acid sequence can be obtained by integrating at least one
5 additional copy of the sequence into the host cell genome or by including an amplifiable selectable marker gene with the nucleic acid sequence where cells containing amplified copies of the selectable marker gene, and thereby additional copies of the nucleic acid sequence, can be selected for by culturing the cells in the presence of the appropriate selectable agent.

The procedures used to ligate the elements described above to construct the recombinant
10 expression vectors of the present invention are well known to one skilled in the art (see, *e.g.*, Sambrook *et al.*, 1989, *supra*).

Host Cells

The present invention also relates to recombinant host cells, comprising a nucleic acid
15 sequence of the invention, which are advantageously used in the recombinant production of the polypeptides. The term "host cell" encompasses any progeny of a parent cell which is not identical to the parent cell due to mutations that occur during replication.

A vector comprising a nucleic acid sequence of the present invention is introduced into a host cell so that the vector is maintained as a chromosomal integrant or as a self-replicating
20 extra-chromosomal vector. Integration is generally considered to be an advantage as the nucleic acid sequence is more likely to be stably maintained in the cell. Integration of the vector into the host chromosome may occur by homologous or non-homologous recombination as described above.

The choice of a host cell will to a large extent depend upon the gene encoding the
25 polypeptide and its source. The host cell may be a unicellular microorganism, *e.g.*, a prokaryote, or a non-unicellular microorganism, *e.g.*, a eukaryote. Useful unicellular cells are bacterial cells such as gram positive bacteria including, but not limited to, a *Bacillus* cell, *e.g.*, *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus brevis*, *Bacillus circulans*, *Bacillus clausii*, *Bacillus coagulans*, *Bacillus lautus*, *Bacillus lentus*, *Bacillus licheniformis*, *Bacillus megaterium*, *Bacillus stearothermophilus*, *Bacillus subtilis*, and *Bacillus thuringiensis*; or a
30 *Streptomyces* cell, *e.g.*, *Streptomyces lividans* or *Streptomyces murinus*, or gram negative bacteria such as *E. coli* and *Pseudomonas* sp. In a preferred embodiment, the bacterial host cell is a *Bacillus lentus*, *Bacillus licheniformis*, *Bacillus stearothermophilus*, or *Bacillus subtilis* cell. The introduction of a vector into a bacterial host cell may, for instance, be effected by
35 protoplast transformation (see, *e.g.*, Chang and Cohen, 1979, *Molecular General Genetics* 168: 111-115), by using competent cells (see, *e.g.*, Young and Spizizin, 1961, *Journal of Bacteriology* 81: 823-829, or Dubnau and Davidoff-Abelson, 1971, *Journal of Molecular Biology* 56: 209-221), by electroporation (see, *e.g.*, Shigekawa and Dower, 1988,

Biotechniques 6: 742-751), or by conjugation (see, *e.g.*, Koehler and Thorne, 1987, *Journal of Bacteriology* 169: 5771-5278).

The host cell may be a eukaryote, such as a mammalian cell, an insect cell, a plant cell or a fungal cell. Useful mammalian cells include Chinese hamster ovary (CHO) cells, HeLa cells, baby hamster kidney (BHK) cells, COS cells, or any number of other immortalized cell lines available, *e.g.*, from the American Type Culture Collection.

In a preferred embodiment, the host cell is a fungal cell. "Fungi" as used herein includes the phyla Ascomycota, Basidiomycota, Chytridiomycota, and Zygomycota (as defined by Hawksworth *et al.*, In, *Ainsworth and Bisby's Dictionary of The Fungi*, 8th edition, 1995, CAB International, University Press, Cambridge, UK) as well as the Oomycota (as cited in Hawksworth *et al.*, 1995, *supra*, page 171) and all mitosporic fungi (Hawksworth *et al.*, 1995, *supra*). Representative groups of Ascomycota include, *e.g.*, *Neurospora*, *Eupenicillium* (= *Penicillium*), *Emmericella* (= *Aspergillus*), *Eurotium* (= *Aspergillus*), and the true yeasts listed below. Examples of Basidiomycota include mushrooms, rusts, and smuts. Representative groups of Chytridiomycota include, *e.g.*, *Allomyces*, *Blastocladiella*, *Coelomomyces*, and aquatic fungi. Representative groups of Oomycota include, *e.g.*, Saprolegniomycetous aquatic fungi (water molds) such as *Achlya*. Examples of mitosporic fungi include *Aspergillus*, *Penicillium*, *Candida*, and *Alternaria*. Representative groups of Zygomycota include, *e.g.*, *Rhizopus* and *Mucor*.

In a more preferred embodiment, the fungal host cell is a yeast cell. "Yeast" as used herein includes ascosporeogenous yeast (Endomycetales), basidiosporogenous yeast, and yeast belonging to the Fungi Imperfecti (Blastomycetes). The ascosporeogenous yeasts are divided into the families Spermophthoraceae and Saccharomycetaceae. The latter is comprised of four subfamilies, Schizosaccharomycoideae (*e.g.*, genus *Schizosaccharomyces*), Nadsonioideae, Lipomycoideae, and Saccharomycoideae (*e.g.*, genera *Kluyveromyces*, *Pichia*, and *Saccharomyces*). The basidiosporogenous yeasts include the genera *Leucosporidium*, *Rhodospiridium*, *Sporidiobolus*, *Filobasidium*, and *Filobasidiella*. Yeast belonging to the Fungi Imperfecti are divided into two families, Sporobolomycetaceae (*e.g.*, genera *Sporobolomyces* and *Bullera*) and Cryptococcaceae (*e.g.*, genus *Candida*). Since the classification of yeast may change in the future, for the purposes of this invention, yeast shall be defined as described in *Biology and Activities of Yeast* (Skinner, F.A., Passmore, S.M., and Davenport, R.R., eds, Soc. App. Bacteriol. Symposium Series No. 9, 1980. The biology of yeast and manipulation of yeast genetics are well known in the art (see, *e.g.*, *Biochemistry and Genetics of Yeast*, Bacil, M., Horecker, B.J., and Stopani, A.O.M., editors, 2nd edition, 1987; *The Yeasts*, Rose, A.H., and Harrison, J.S., editors, 2nd edition, 1987; and *The Molecular Biology of the Yeast Saccharomyces*, Strathern *et al.*, editors, 1981).

In an even more preferred embodiment, the yeast host cell is a cell of a species of *Candida*, *Hansenula*, *Kluyveromyces*, *Pichia*, *Saccharomyces*, *Schizosaccharomyces*, or

Yarrowia.

In a most preferred embodiment, the yeast host cell is a *Saccharomyces carlsbergensis*, *Saccharomyces cerevisiae*, *Saccharomyces diastaticus*, *Saccharomyces douglasii*, *Saccharomyces kluyveri*, *Saccharomyces norbensis* or *Saccharomyces oviformis* cell. In another most preferred embodiment, the yeast host cell is a *Kluyveromyces lactis* cell. In another most preferred embodiment, the yeast host cell is a *Yarrowia lipolytica* cell.

In another more preferred embodiment, the fungal host cell is a filamentous fungal cell. "Filamentous fungi" include all filamentous forms of the subdivision Eumycota and Oomycota (as defined by Hawksworth *et al.*, 1995, *supra*). The filamentous fungi are characterized by a mycelial wall composed of chitin, cellulose, glucan, chitosan, mannan, and other complex polysaccharides. Vegetative growth is by hyphal elongation and carbon catabolism is obligately aerobic. In contrast, vegetative growth by yeasts such as *Saccharomyces cerevisiae* is by budding of a unicellular thallus and carbon catabolism may be fermentative. In a more preferred embodiment, the filamentous fungal host cell is a cell of a species of, but not limited to, *Acremonium*, *Aspergillus*, *Fusarium*, *Humicola*, *Mucor*, *Myceliophthora*, *Neurospora*, *Penicillium*, *Thielavia*, *Tolypocladium*, and *Trichoderma*.

In an even more preferred embodiment, the filamentous fungal host cell is an *Aspergillus* cell. In another even more preferred embodiment, the filamentous fungal host cell is an *Acremonium* cell. In another even more preferred embodiment, the filamentous fungal host cell is a *Fusarium* cell. In another even more preferred embodiment, the filamentous fungal host cell is a *Humicola* cell. In another even more preferred embodiment, the filamentous fungal host cell is a *Mucor* cell. In another even more preferred embodiment, the filamentous fungal host cell is a *Myceliophthora* cell. In another even more preferred embodiment, the filamentous fungal host cell is a *Neurospora* cell. In another even more preferred embodiment, the filamentous fungal host cell is a *Penicillium* cell. In another even more preferred embodiment, the filamentous fungal host cell is a *Thielavia* cell. In another even more preferred embodiment, the filamentous fungal host cell is a *Tolypocladium* cell. In another even more preferred embodiment, the filamentous fungal host cell is a *Trichoderma* cell.

In a most preferred embodiment, the filamentous fungal host cell is an *Aspergillus awamori*, *Aspergillus foetidus*, *Aspergillus japonicus*, *Aspergillus nidulans*, *Aspergillus niger*, or *Aspergillus oryzae* cell. In another most preferred embodiment, the filamentous fungal host cell is a *Fusarium bactridioides*, *Fusarium cerealis*, *Fusarium crookwellense*, *Fusarium culmorum*, *Fusarium graminearum*, *Fusarium graminum*, *Fusarium heterosporum*, *Fusarium negundi*, *Fusarium oxysporum*, *Fusarium reticulatum*, *Fusarium roseum*, *Fusarium sambucinum*, *Fusarium sarcochroum*, *Fusarium solani*, *Fusarium sporotrichioides*, *Fusarium sulphureum*, *Fusarium torulosum*, *Fusarium trichothecioides*, or *Fusarium venenatum* cell. In an even most preferred embodiment, the filamentous fungal parent cell is a *Fusarium*

venenatum (Nirenberg sp. nov.) cell. In another most preferred embodiment, the filamentous fungal host cell is a *Humicola insolens* or *Humicola lanuginosa* cell. In another most preferred embodiment, the filamentous fungal host cell is a *Mucor miehei* cell. In another most preferred embodiment, the filamentous fungal host cell is a *Myceliophthora thermophilum* cell. In another most preferred embodiment, the filamentous fungal host cell is a *Neurospora crassa* cell. In another most preferred embodiment, the filamentous fungal host cell is a *Penicillium purpurogenum* cell. In another most preferred embodiment, the filamentous fungal host cell is a *Thielavia terrestris* cell. In another most preferred embodiment, the *Trichoderma* cell is a *Trichoderma harzianum*, *Trichoderma koningii*, *Trichoderma longibrachiatum*, *Trichoderma reesei* or *Trichoderma viride* cell.

Fungal cells may be transformed by a process involving protoplast formation, transformation of the protoplasts, and regeneration of the cell wall in a manner known *per se*. Suitable procedures for transformation of *Aspergillus* host cells are described in EP 238 023 and Yelton *et al.*, 1984, *Proceedings of the National Academy of Sciences USA* 81: 1470-1474. Suitable methods for transforming *Fusarium* species are described by Malardier *et al.*, 1989, *Gene* 78: 147-156 and WO 96/00787. Yeast may be transformed using the procedures described by Becker and Guarente, In Abelson, J.N. and Simon, M.I., editors, *Guide to Yeast Genetics and Molecular Biology*, Methods in Enzymology, Volume 194, pp 182-187, Academic Press, Inc., New York; Ito *et al.*, 1983, *Journal of Bacteriology* 153: 163; and Hinnen *et al.*, 1978, *Proceedings of the National Academy of Sciences USA* 75: 1920. Mammalian cells may be transformed by direct uptake using the calcium phosphate precipitation method of Graham and Van der Eb (1978, *Virology* 52: 546).

Methods of Production

The present invention also relates to methods for producing a polypeptide of the present invention comprising (a) cultivating a strain, which in its wild-type form is capable of producing the polypeptide, to produce a supernatant comprising the polypeptide; and (b) recovering the polypeptide. Preferably, the strain is of the genus *Aspergillus*.

The present invention also relates to methods for producing a polypeptide of the present invention comprising (a) cultivating a host cell under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.

The present invention further relates to methods for producing a polypeptide of the present invention comprising (a) cultivating a homologously recombinant cell, having incorporated therein a new transcription unit comprising a regulatory sequence, an exon, and/or a splice donor site operably linked to a second exon of an endogenous nucleic acid sequence encoding the polypeptide, under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide. The methods are based on the use of gene activation technology, for example, as described in U.S. Patent No. 5,641,670. Gene activation

technology is based on activating a gene which is normally unexpressed in a cell or increasing expression of a gene which is expressed at very low levels in a cell. Gene activation technology includes methods of inserting an exogenous DNA construct containing a regulatory sequence, an exon, and/or a splice donor site into the genomic DNA of a cell in such a manner
5 that the insertion results in the production of a new transcription unit in which the regulatory sequence, the exon, and/or the splice donor site are operably linked to and activate expression of the endogenous gene.

In the production methods of the present invention, the cells are cultivated in a nutrient medium suitable for production of the polypeptide using methods known in the art. For
10 example, the cell may be cultivated by shake flask cultivation, small-scale or large-scale fermentation (including continuous, batch, fed-batch, or solid state fermentations) in laboratory or industrial fermentors performed in a suitable medium and under conditions allowing the polypeptide to be expressed and/or isolated. The cultivation takes place in a suitable nutrient medium comprising carbon and nitrogen sources and inorganic salts, using procedures known
15 in the art (see, *e.g.*, references for bacteria and yeast; Bennett, J.W. and LaSure, L., editors, *More Gene Manipulations in Fungi*, Academic Press, CA, 1991). Suitable media are available from commercial suppliers or may be prepared according to published compositions (*e.g.*, in catalogues of the American Type Culture Collection). If the polypeptide is secreted into the nutrient medium, the polypeptide can be recovered directly from the medium. If the
20 polypeptide is not secreted, it can be recovered from cell lysates.

The polypeptides may be detected using methods known in the art that are specific for the polypeptides. These detection methods may include use of specific antibodies, formation of an enzyme product, or disappearance of an enzyme substrate. For example, an enzyme assay may be used to determine the activity of the polypeptide. Procedures for determining dipeptidyl
25 aminopeptidase activity are known in the art and include, *e.g.*, measuring the initial rate of hydrolysis of a *p*-nitroanilide at 405 nm.

The resulting polypeptide may be recovered by methods known in the art. For example, the polypeptide may be recovered from the nutrient medium by conventional procedures including, but not limited to, centrifugation, filtration, extraction, spray-drying,
30 evaporation, or precipitation.

The polypeptides of the present invention may be purified by a variety of procedures known in the art including, but not limited to, chromatography (*e.g.*, ion exchange, affinity, hydrophobic, chromatofocusing, and size exclusion), electrophoretic procedures (*e.g.*, preparative isoelectric focusing, differential solubility (*e.g.*, ammonium sulfate precipitation),
35 SDS-PAGE, or extraction (see, *e.g.*, *Protein Purification*, J.-C. Janson and Lars Ryden, editors, VCH Publishers, New York, 1989).

Removal or Reduction of Dipeptidyl Aminopeptidase Activity

The present invention also relates to methods for producing a mutant cell of a parent cell, which comprises disrupting or deleting a nucleic acid sequence encoding the polypeptide or a control sequence thereof, which results in the mutant cell producing less of the polypeptide than the parent cell.

5 The construction of strains which have reduced dipeptidyl aminopeptidase activity may be conveniently accomplished by modification or inactivation of a nucleic acid sequence necessary for expression of the polypeptide having dipeptidyl aminopeptidase activity in the cell. The nucleic acid sequence to be modified or inactivated may be, for example, a nucleic acid sequence encoding the polypeptide or a part thereof essential for exhibiting dipeptidyl
10 aminopeptidase activity, or the nucleic acid sequence may have a regulatory function required for the expression of the polypeptide from the coding sequence of the nucleic acid sequence. An example of such a regulatory or control sequence may be a promoter sequence or a functional part thereof, *i.e.*, a part which is sufficient for affecting expression of the polypeptide. Other control sequences for possible modification include, but are not limited to,
15 a leader, a polyadenylation sequence, a propeptide sequence, a signal sequence, and a termination terminator.

Modification or inactivation of the nucleic acid sequence may be performed by subjecting the cell to mutagenesis and selecting for cells in which the dipeptidyl aminopeptidase producing capability has been reduced or eliminated. The mutagenesis, which may be specific
20 or random, may be performed, for example, by use of a suitable physical or chemical mutagenizing agent, by use of a suitable oligonucleotide, or by subjecting the DNA sequence to PCR generated mutagenesis. Furthermore, the mutagenesis may be performed by use of any combination of these mutagenizing agents.

Examples of a physical or chemical mutagenizing agent suitable for the present purpose
25 include ultraviolet (UV) irradiation, hydroxylamine, N-methyl-N'-nitro-N-nitrosoguanidine (MNNG), O-methyl hydroxylamine, nitrous acid, ethyl methane sulphonate (EMS), sodium bisulphite, formic acid, and nucleotide analogues.

When such agents are used, the mutagenesis is typically performed by incubating the cell to be mutagenized in the presence of the mutagenizing agent of choice under suitable
30 conditions, and selecting for cells exhibiting reduced or no expression of dipeptidyl aminopeptidase activity.

Modification or inactivation of production of a polypeptide of the present invention may be accomplished by introduction, substitution, or removal of one or more nucleotides in the nucleic acid sequence encoding the polypeptide or a regulatory element required for the
35 transcription or translation thereof. For example, nucleotides may be inserted or removed so as to result in the introduction of a stop codon, the removal of the start codon, or a change of the open reading frame. Such modification or inactivation may be accomplished by site-directed mutagenesis or PCR generated mutagenesis in accordance with methods known in the art.

Although, in principle, the modification may be performed *in vivo*, *i.e.*, directly on the cell expressing the nucleic acid sequence to be modified, it is preferred that the modification be performed *in vitro* as exemplified below.

5 An example of a convenient way to inactivate or reduce production by a host cell of choice is based on techniques of gene replacement or gene interruption. For example, in the gene interruption method, a nucleic acid sequence corresponding to the endogenous gene or gene fragment of interest is mutagenized *in vitro* to produce a defective nucleic acid sequence which is then transformed into the host cell to produce a defective gene. By homologous recombination, the defective nucleic acid sequence replaces the endogenous gene or gene
10 fragment. It may be desirable that the defective gene or gene fragment also encodes a marker which may be used for selection of transformants in which the gene encoding the polypeptide has been modified or destroyed.

Alternatively, modification or inactivation of the nucleic acid sequence encoding a polypeptide of the present invention may be performed by established anti-sense techniques
15 using a nucleotide sequence complementary to the polypeptide encoding sequence. More specifically, production of the polypeptide by a cell may be reduced or eliminated by introducing a nucleotide sequence complementary to the nucleic acid sequence encoding the polypeptide which may be transcribed in the cell and is capable of hybridizing to the polypeptide mRNA produced in the cell. Under conditions allowing the complementary anti-
20 sense nucleotide sequence to hybridize to the polypeptide mRNA, the amount of polypeptide translated is thus reduced or eliminated.

It is preferred that the cell to be modified in accordance with the methods of the present invention is of microbial origin, for example, a fungal strain which is suitable for the production of desired protein products, either homologous or heterologous to the cell.

25 The present invention further relates to a mutant cell of a parent cell which comprises a disruption or deletion of a nucleic acid sequence encoding the polypeptide or a control sequence thereof, which results in the mutant cell producing less of the polypeptide than the parent cell.

The polypeptide-deficient mutant cells so created are particularly useful as host cells for the expression of homologous and/or heterologous polypeptides. Therefore, the present
30 invention further relates to methods for producing a homologous or heterologous polypeptide comprising (a) culturing the mutant cell under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide. In the present context, the term "heterologous polypeptides" is defined herein as polypeptides which are not native to the host cell, a native protein in which modifications have been made to alter the native sequence, or a native protein
35 whose expression is quantitatively altered as a result of a manipulation of the host cell by recombinant DNA techniques.

In a still further aspect, the present invention relates to a method for producing a protein product essentially free of dipeptidyl aminopeptidase activity by fermentation of a cell which

produces both a polypeptide of the present invention as well as the protein product of interest. The method comprises adding an effective amount of an agent capable of inhibiting dipeptidyl aminopeptidase activity to the fermentation broth either during or after the fermentation has been completed, recovering the product of interest from the fermentation broth, and optionally
5 subjecting the recovered product to further purification.

In a still further alternative aspect, the present invention relates to a method for producing a protein product essentially free of dipeptidyl aminopeptidase activity, wherein the protein product of interest is encoded by a DNA sequence present in a cell encoding a polypeptide of the present invention. The method comprises cultivating the cell under
10 conditions permitting the expression of the product, subjecting the resultant culture broth to a combined pH and temperature treatment so as to reduce the dipeptidyl aminopeptidase activity substantially, and recovering the product from the culture broth. Alternatively, the combined pH and temperature treatment may be performed on an enzyme preparation recovered from the culture broth. The combined pH and temperature treatment may optionally be used in
15 combination with a treatment with a dipeptidyl aminopeptidase inhibitor.

The combined pH and temperature treatment is preferably carried out at a pH in the range of 9-11 and a temperature in the range of 40-75°C for a sufficient period of time to attain the desired effect, where typically, 30 to 60 minutes is sufficient.

In accordance with this aspect of the invention, it is possible to remove at least 60%,
20 preferably at least 75%, more preferably at least 85%, still more preferably at least 95%, and most preferably at least 99% of the dipeptidyl aminopeptidase activity. It is contemplated that a complete removal of dipeptidyl aminopeptidase activity may be obtained by use of these methods.

The methods used for cultivation and purification of the product of interest may be
25 performed by methods known in the art.

The methods of the present invention for producing an essentially dipeptidyl aminopeptidase-free product is of particular interest in the production of eukaryotic polypeptides, in particular fungal proteins such as enzymes. The enzyme may be selected from, *e.g.*, an amylolytic enzyme, a lipolytic enzyme, a proteolytic enzyme, a cellulytic
30 enzyme, an oxidoreductase or a plant cell-wall degrading enzyme. Examples of such enzymes include an aminopeptidase, an amylase, an amyloglucosidase, a carbohydrase, a carboxypeptidase, a catalase, a cellulase, a chitinase, a cutinase, a cyclodextrin glycosyltransferase, a deoxyribonuclease, an esterase, a galactosidase, a beta-galactosidase, a glucoamylase, a glucose oxidase, a glucosidase, a haloperoxidase, a hemicellulase, an
35 invertase, an isomerase, a laccase, a ligase, a lipase, a lyase, a mannosidase, an oxidase, a pectinolytic enzyme, a peroxidase, a phytase, a phenoloxidase, a polyphenoloxidase, a proteolytic enzyme, a ribonuclease, a transferase, a transglutaminase, or a xylanase. The dipeptidyl aminopeptidase-deficient cells may also be used to express heterologous proteins of

pharmaceutical interest such as hormones, growth factors, receptors, and the like.

It will be understood that the term "eukaryotic polypeptides" includes not only native polypeptides, but also those polypeptides, *e.g.*, enzymes, which have been modified by amino acid substitutions, deletions or additions, or other such modifications to enhance activity, thermostability, pH tolerance and the like.

In a further aspect, the present invention relates to a protein product essentially free from dipeptidyl aminopeptidase activity which is produced by a method of the present invention.

Methods of Producing Protein Hydrolysates

The polypeptides of the present invention may be used in the production of protein hydrolysates for enhancing the degree of hydrolysis and flavor development.

The present invention further relates to methods for using a polypeptide of the present invention in combination with an endopeptidase to produce a high degree of hydrolysis of a protein-rich material. The method comprises treating of a proteinaceous substrate with the polypeptide and an endopeptidase. The substrate may be treated with the enzymes concurrently or consecutively.

A polypeptide of the present invention is added to the proteinaceous substrate in an effective amount conventionally employed in protein hydrolysis processes, preferably in the range of from about 0.1 to about 100,000 dipeptidyl aminopeptidase units (DPAPU) per 100 g of protein, and more preferably in the range of from about 1 to about 10,000 dipeptidyl aminopeptidase units per 100 g of protein. As defined herein, one dipeptidyl aminopeptidase Unit (DPAPU) is the amount of enzyme needed to release 1 micromole of p-nitroanilide per minute from Ala-Pro-p-nitroanilide (Sigma Chemical Co., St. Louis MO) under specified conditions.

The endopeptidase may be obtained from a strain of *Bacillus*, preferably *Bacillus licheniformis* or *Bacillus subtilis*, a strain of *Staphylococcus*, preferably *Staphylococcus aureus*, a strain of *Streptomyces*, preferably *Streptomyces thermovularis* or *Streptomyces griseus*, a strain of *Actinomyces* species, a strain of *Aspergillus*, preferably *Aspergillus aculeatus*, *Aspergillus awamori*, *Aspergillus foetidus*, *Aspergillus nidulans*, *Aspergillus niger*, or *Aspergillus oryzae*, or a strain of *Fusarium*, preferably *Fusarium venenatum*.

The endopeptidase is added to the proteinaceous substrate in an effective amount conventionally employed in protein hydrolysis processes, preferably in the range of from about 0.05 to about 15 AU/100 g of protein, and more preferably from about 0.1 to about 8 AU/100 g of protein. One AU (Anson Unit) is defined as the amount of enzyme which under standard conditions (*i.e.*, 25°C, pH 7.5 and 10 min. reaction time) digests hemoglobin at an initial rate such that there is liberated per minute an amount of TCA soluble product which gives the same color with phenol reagent as one milli-equivalent of tyrosine. The analytical method AF 4/5 is

available upon request from Novo Nordisk A/S, Denmark, which is incorporated herein by reference.

The enzymatic treatment, *i.e.*, the incubation of the substrate with the enzyme preparations, may take place at any convenient temperature at which the enzyme preparation does not become inactivated, preferably in the range of from about 20°C to about 70°C. In accordance with established practice, the enzyme preparations may be suitably inactivated by increasing the temperature of the incubation mixture to a temperature where the enzymes become inactivated, *e.g.*, to above about 70°C, or similarly by decreasing the pH of the incubation mixture to a point where the enzymes become inactivated, *e.g.*, below about 4.0.

Furthermore, the methods of the present invention result in enhancement of the degree of hydrolysis of a proteinaceous substrate. As used herein, the degree of hydrolysis (DH) is the percentage of the total number of amino bonds in a protein that has been hydrolyzed by a proteolytic enzyme.

In another aspect of the present invention, the hydrolysates have an increased content of Ala, Arg, Asp, Gly, and/or Val, *e.g.*, 1.1 times greater.

In another aspect of the present invention, a polypeptide of the present invention acts synergistically with an aminopeptidase to hydrolyze a tripeptide where either enzyme activity alone does not hydrolyze the tripeptide. In a preferred embodiment, the aminopeptidase is aminopeptidase I obtained from *Aspergillus oryzae* as described in WO 96/28542.

The present invention also relates to methods for obtaining a protein hydrolysate enriched in free glutamic acid and/or peptide bound glutamic acid residues, which method comprises:

(a) subjecting the substrate to a deamidation process; and

(b) subjecting the substrate to the action of a polypeptide having dipeptidyl aminopeptidase activity.

The two steps may be performed simultaneously, or the second step may be performed subsequent to the first step.

These methods of the present invention produce protein hydrolysates of excellent flavor because glutamic acid (Glu), whether free or peptide bound, plays an important role in the flavor and palatability of protein hydrolysates. These method also produce protein hydrolysates having improved functionality, in particular, improved solubility, improved emulsifying properties, increased degree of hydrolysis, and improved foaming properties.

The conversion of amides (glutamine or asparagine) into charged acids (glutamic acid or aspartic acid) via the liberation of ammonia is known as deamidation. Deamidation may take place as a non-enzymatic or as an enzymatic deamidation process.

In a preferred embodiment, the deamidation is carried out as an enzymatic deamidation process, *e.g.*, by subjecting the substrate to a transglutaminase and/or peptidoglutaminase.

The transglutaminase may be of any convenient source including mammals, see *e.g.*,

JP 1050382 and JP 5023182, including activated Factor XIII, see *e.g.*, WO 93/15234; those derived from fish, see *e.g.*, EP 555,649; and those obtained from microorganisms, see *e.g.*, EP 379,606, WO 96/06931 and WO 96/22366. In a preferred embodiment, the transglutaminase is obtained from an Oomycete, including a strain of *Phytophthora*, preferably *Phytophthora cactorum*, or a strain of *Pythium*, preferably *Pythium irregulare*, *Pythium* sp., *Pythium intermedium*, *Pythium ultimum*, or *Pythium peritum* (or *Pythium periplocum*). In another preferred embodiment, the transglutaminase is of bacterial origin and is obtained from a strain of *Bacillus*, preferably *Bacillus subtilis*, a strain of *Streptovorticillium*, preferably *Streptovorticillium mobaraensis*, *Streptovorticillium griseocarneum*, or *Streptovorticillium cinnamoneum*, and a strain of *Streptomyces*, preferably *Streptomyces lydicus*.

The peptidoglutaminase may be a peptidoglutaminase I (peptidyl-glutaminase; EC 3.5.1.43), or a peptidoglutaminase II (protein-glutamine glutaminase; EC 3.5.1.44), or any mixture thereof. The peptidoglutaminase may be obtained from a strain of *Aspergillus*, preferably *Aspergillus japonicus*, a strain of *Bacillus*, preferably *Bacillus circulans*, a strain of *Cryptococcus*, preferably *Cryptococcus albidus*, or a strain of *Debaryomyces*, preferably *Debaryomyces hansenii*.

The transglutaminase is added to the proteinaceous substrate in an effective amount conventionally employed in deamidation processes, preferably in the range of from about 0.01 to about 5% (w/w), and more preferably in the range of from about 0.1 to about 1% (w/w) of enzyme preparation relating to the amount of substrate.

The peptidoglutaminase is added to the proteinaceous substrate in an effective amount conventionally employed in deamidation processes, preferably in the range of from about 0.01 to about 100,000 PGase Units per 100 g of substrate, and more preferably in the range of from about 0.1 to about 10,000 PGase Units per 100 g of substrate.

The peptidoglutaminase activity may be determined according to the procedure of Cedrangoro *et al.* (1965, *Enzymologia* 29: 143). According to this procedure, 0.5 ml of an enzyme sample, adjusted to pH 6.5 with 1 N NaOH, is charged into a small vessel. Then 1 ml of a borate pH 10.8 buffer solution is added to the vessel. The discharged ammonia is absorbed by 5 N sulphuric acid, and by use of Nessler's reagent the mixture is allowed to form color which is measured at 420 nm. One PGase unit is the amount of enzyme capable of producing 1 micromole of ammonia per minute under these conditions.

Alternatively, the peptidoglutaminase activity may be determined according to the procedure described in US 3,857,967 or Example 17 below.

In step (b) of the methods of the present invention, the substrate is subjected to a polypeptide of the present invention. A polypeptide of the present invention is added to the proteinaceous substrate in an effective amount conventionally employed in protein hydrolysis processes, preferably in the range of from about 0.001 to about 0.5 AU/100 g of substrate, more preferably in the range of from about 0.01 to about 0.1 AU/100 g of substrate.

In another embodiment, the methods of the present invention may be used to produce a hydrolysate enriched in free glutamic acid and/or peptide bound glutamic acid residues further comprise:

(c) subjecting the substrate to one or more unspecific acting endo- and/or exo-peptidase enzymes.

This step may take place simultaneously with steps (a) and (b), or may follow steps (a) and (b).

In a preferred embodiment, the unspecific acting endo- and/or exo-peptidase enzyme is obtained from a strain of *Aspergillus*, preferably *Aspergillus niger*, *Aspergillus oryzae*, or *Aspergillus sojae*, or a strain of *Bacillus*, preferably *Bacillus amyloliquefaciens*, *Bacillus lentus*, *Bacillus licheniformis*, or *Bacillus subtilis*.

The unspecific acting endo- and/or exo-peptidase enzyme is added to the substrate in an effective amount conventionally employed in protein hydrolysis processes, preferably in the range of from about 0.05 to about 15 CPU/100 g of substrate, and more preferably in the range of from about 0.1 to about 5 CPU/100 g of substrate. One CPU (Casein Protease Unit) is defined as the amount of enzyme liberating 1 micromole of primary amino groups (determined by comparison with a serine standard) per minute from casein under standard conditions, *i.e.*, incubation for 30 minutes at 25°C and pH 9.5. The analytical method AF 228/1, which is incorporated herein by reference, is available upon request from Novo Nordisk A/S, Bagsværd, Denmark.

Each enzymatic treatment may take place at any temperature at which the enzyme preparation does not become inactivated, preferably in the range of from about 20°C to about 70°C. The enzyme preparation may then be inactivated by increasing the temperature, *e.g.*, to above about 70°C, or by decreasing the pH, *e.g.*, below about 4.0.

The proteinaceous substrate used in the methods of the present invention may consist of intact proteins, prehydrolyzed proteins (*i.e.*, peptides), or a mixture thereof. The proteinaceous substrate may be of vegetable or animal origin. Preferably, the proteinaceous substrate is of vegetable origin, *e.g.*, soy protein, grain protein, *e.g.*, wheat gluten, corn gluten, barley, rye, oat, rice, zein, lupine, cotton seed protein, rape seed protein, peanut, alfalfa protein, pea protein, fabaceous bean protein, sesame seed protein, or sunflower. A proteinaceous substrate of animal origin may be whey protein, casein, meat proteins, fish protein, red blood cells, egg white, gelatin, or lactoalbumin.

The present invention also relates to protein hydrolysates produced by these methods.

Other Uses

The present invention also relates to methods of deactivating enzymes with a polypeptide of the present invention.

Furthermore, a polypeptide of the present invention may be useful for a number of

purposes in which a specific cleavage of peptide sequences is desirable. For instance, some proteins or peptides are synthesized in the form of inactive precursors comprising a number of additional amino acid residues at the N-terminal of the mature protein. A polypeptide of the present invention could provide the necessary post-translational processing to activate such precursor proteins.

Compositions

In a still further aspect, the present invention relates to polypeptide compositions comprising a polypeptide of the present invention. Preferably, the compositions are enriched in a polypeptide of the present invention. In the present context, the term "enriched" indicates that the dipeptidyl aminopeptidase activity of the polypeptide composition has been increased, *e.g.*, with an enrichment factor of 1.1.

The polypeptide composition may comprise a polypeptide of the invention as the major enzymatic component, *e.g.*, a mono-component polypeptide composition. Alternatively, the composition may comprise multiple enzymatic activities, such as an aminopeptidase, an amylase, a carbohydrase, a carboxypeptidase, a catalase, a cellulase, a chitinase, a cutinase, a cyclodextrin glycosyltransferase, a deoxyribonuclease, an esterase, an alpha-galactosidase, a beta-galactosidase, a glucoamylase, an alpha-glucosidase, a beta-glucosidase, a haloperoxidase, an invertase, a laccase, a lipase, a mannosidase, an oxidase, a pectinolytic enzyme, a peptidoglutaminase, a peroxidase, a phytase, a polyphenoloxidase, a proteolytic enzyme, a ribonuclease, a transglutaminase, or a xylanase. The additional enzyme(s) may be producible by means of a microorganism belonging to the genus *Aspergillus*, preferably *Aspergillus aculeatus*, *Aspergillus awamori*, *Aspergillus niger*, or *Aspergillus oryzae*, or *Trichoderma*, *Humicola*, preferably *Humicola insolens*, or *Fusarium*, preferably *Fusarium bac-tridioides*, *Fusarium cerealis*, *Fusarium crookwellense*, *Fusarium culmorum*, *Fusarium graminearum*, *Fusarium graminum*, *Fusarium heterosporum*, *Fusarium negundi*, *Fusarium oxysporum*, *Fusarium reticulatum*, *Fusarium roseum*, *Fusarium sambucinum*, *Fusarium sarcochroum*, *Fusarium sporotrichioides*, *Fusarium sulphureum*, *Fusarium torulosum*, *Fusarium trichothecioides*, or *Fusarium venenatum*.

In a preferred embodiment, the invention relates to a flavor-improving composition comprising a polypeptide with dipeptidyl aminopeptidase activity and a carrier. Any suitable carrier known in the art may be used. In another preferred embodiment, the flavor-improving composition further comprises an endopeptidase. In another preferred embodiment, the flavoring composition further comprises one or more unspecific-acting endo- and/or exo-peptidase enzymes. In another preferred embodiment, the flavoring composition further comprises one or more specific-acting endo- and/or exo-peptidase enzymes.

In a preferred embodiment, the specific acting proteolytic enzyme is an endopeptidase such as a glutamyl endopeptidase (EC 3.4.21.19); a lysyl endopeptidase (EC 3.4.21.50); a

leucyl endopeptidase (EC 3.4.21.57); a glycy endopeptidase (EC 3.4.22.25); a prolyl endopeptidase (EC 3.4.21.26); trypsin (EC 3.4.21.4) or a trypsin-like (lysine/arginine specific) endopeptidase; or a peptidyl-Asp metalloendopeptidase (EC 3.4.24.33).

The glutamyl endopeptidase (EC 3.4.21.19) may preferably be obtained from a *Bacillus* strain, in particular *Bacillus licheniformis* and *Bacillus subtilis*, a *Staphylococcus* strain, in particular *Staphylococcus aureus*, a *Streptomyces* strain, in particular *Streptomyces thermovulgaris* and *Streptomyces griseus*, or a *Actinomyces* strain.

The lysyl endopeptidase (EC 3.4.21.50) may preferably be obtained from a *Achromobacter* strain, in particular *Achromobacter lyticus*, a *Lysobacter* strain, in particular *Lysobacter enzymogenes*, or a *Pseudomonas* strain, in particular *Pseudomonas aeruginosa*.

The leucyl endopeptidase (EC 3.4.21.57) may be of plant origin.

The glycy endopeptidase (EC 3.4.22.25) may preferably be obtained from the papaya plant (*Carica papaya*).

The prolyl endopeptidase (EC 3.4.21.26) may preferably be obtained from a *Flavobacterium* strain, or it may be of plant origin.

The trypsin-like endopeptidase may preferably be obtained from a *Fusarium* strain, in particular *Fusarium oxysporum*, *e.g.*, as described in WO 89/06270 or WO 94/25583.

The peptidyl-Asp metalloendopeptidase (EC 3.4.24.33) may preferably be obtained from a *Pseudomonas* strain, in particular *Pseudomonas fragi*.

In another preferred embodiment, the specific acting proteolytic enzyme is an exopeptidase that may act from either end of the peptide.

In a preferred embodiment, the specific acting proteolytic enzyme is an aminopeptidase such as a leucyl aminopeptidase (EC 3.4.11.1); or a tripeptide aminopeptidase (EC 3.4.11.4).

In another preferred embodiment, the specific acting proteolytic enzyme is a carboxypeptidase such as a proline carboxypeptidase (EC 3.4.16.2); a carboxypeptidase A (EC 3.4.17.1); a carboxypeptidase B (EC 3.4.17.2); a carboxypeptidase C (EC 3.4.16.5); a carboxypeptidase D (EC 3.4.16.6); a lysine (arginine) carboxypeptidase (EC 3.4.17.3); a glycine carboxypeptidase (EC 3.4.17.4); an alanine carboxypeptidase (EC 3.4.17.6); a glutamate carboxypeptidase (EC 3.4.17.11); a peptidyl-dipeptidase A (EC 3.4.15.1); or a peptidyl-dipeptidase (EC 3.4.15.5).

The polypeptide compositions may be prepared in accordance with methods known in the art and may be in the form of a liquid or a dry composition. The polypeptide may be stabilized by methods known in the art.

The present invention also relates to food products, *e.g.*, baked products, comprising a protein hydrolysate obtained by the methods of the present invention. Such food products exhibit enhanced organoleptic qualities, such as improvement in flavor, palatability, mouth feel, aroma and crust color.

In the present context, the term "baked products" includes any food prepared from

dough, either of a soft or a crisp character. Examples of baked products, whether of a white, light or dark type, which may be advantageously produced by the present invention, are bread, in particular white, whole-meal or rye bread, typically in the form of loaves or rolls; French baguette-type breads; pita breads; tacos; cakes; pancakes; biscuits; crisp breads; and the like.

5 Such baked products are conventionally prepared from a dough which comprises flour and water, and which is typically leavened. The dough may be leavened in various ways, such as by adding sodium bicarbonate or the like, or by adding a leaven (fermenting dough), but the dough is preferably leavened by adding a suitable yeast culture such as a culture of *Saccharomyces cerevisiae* (baker's yeast). Any of the commercially available *Saccharomyces*
10 *cerevisiae* strains may be employed.

Further, the dough used in the preparation of the baked products may be fresh or frozen. The preparation of frozen dough is described by K. Kulp and K. Lorenz in "Frozen and Refrigerated Doughs and Batters". A flavor improving composition of the present invention is typically included in the dough in an amount in the range of 0.01-5%, more
15 preferably 0.1-3%.

In the methods of the present invention, a polypeptide of the present invention, an endopeptidase, a transglutaminase, a peptidoglutaminase, one or more specific and/or unspecific acting endo- and/or exo-peptidase enzymes, and/or one or more enzymes specified above may be added, either separately or concurrently, to the mixture from which the dough is
20 made or to any ingredient, *e.g.*, flour, from which the dough is to be made.

The present invention further relates to a pre-mix, *e.g.*, in the form of a flour composition, for dough and/or baked products made from dough, wherein the pre-mix comprises a polypeptide or a flavor-improving composition of the invention and a carrier or baking ingredient, and optionally one or more other enzymes specified above.

25 In another embodiment, the pre-mix comprises a hydrolysate obtained by the methods of the invention.

The pre-mix may be prepared by mixing the relevant enzymes with a suitable carrier such as flour, starch, a sugar or a salt. The pre-mix may contain other dough-improving and/or bread-improving additives.

30 In the present context, the term "pre-mix" is a mixture of baking agents, normally including flour, which has been prepared to permit storage under designated conditions and provide convenience in handling during dough preparation processes. Such a pre-mix may be of advantageous use in industrial and commercial bread-baking plants and facilities, as well as in retail bakeries.

35 The present invention also relates to the use of a hydrolysate produced by the methods of the invention as an additive to food products, such as baked foods, to enhance organoleptic qualities, such as flavor, palatability and aroma.

The hydrolysates enriched in free glutamic acid and/or peptide bound glutamic acid

residues obtained by the methods of the present invention may be used in various industrial applications, in particular, where there is a need for the incorporation of functional proteins.

For example, the present invention also relates to food products comprising a hydrolysate enriched in free glutamic acid and/or peptide bound glutamic acid residues obtained by the method of the invention and to animal feed additives comprising a hydrolysate enriched in free glutamic acid and/or peptide bound glutamic acid residues obtained by the methods of the present invention.

The present invention is further described by the following examples which should not be construed as limiting the scope of the invention.

Examples

Example 1: Purification of FLAVOURZYME™ dipeptidyl aminopeptidase I

Dipeptidyl aminopeptidase I was purified from a FLAVOURZYME™ broth (Novo Nordisk A/S, Bagsværd, Denmark). The FLAVOURZYME™ broth was produced by cultivation of *Aspergillus oryzae* strain 1568 (ATCC 20386) in a medium comprised of carbon and nitrogen sources and trace metals. First, the broth (20 ml containing 720 mg of protein) was diluted with 180 ml of 20 mM sodium phosphate pH 7.0 buffer and filtered using Nalgene Filterware equipped with a 0.45 µm filter. The filtered solution was loaded onto a 24 x 130 mm column containing 31 ml of Q-Sepharose, Big Beads (Pharmacia Biotech AB, Uppsala, Sweden) pre-equilibrated with 20 mM sodium phosphate pH 7.0 buffer. The dipeptidyl aminopeptidase I was eluted using 20 mM sodium acetate buffer, pH 3.5. Dipeptidyl aminopeptidase I activity was monitored at 405 nm using 1 mg/ml Ala-Pro-*p*-nitroanilide as substrate in 50 mM sodium phosphate pH 7.5 buffer. The resultant solution containing dipeptidyl aminopeptidase I activity was concentrated to 38 ml by ultrafiltration with a PM10 membrane (Amicon, New Bedford, MA) and then the pH was adjusted to pH 7.0 using a 20 mM Na₂HPO₄ solution.

The resultant solution was loaded onto a 24 x 130 mm column containing 31 ml of Q-Sepharose Big Beads (Pharmacia Biotech AB, Uppsala, Sweden) pre-equilibrated with 20 mM sodium phosphate pH 7.0 buffer. The dipeptidyl aminopeptidase I was eluted with a 0 to 0.3 M NaCl gradient in 20 mM sodium phosphate pH 7.0 buffer. Fractions were monitored for dipeptidyl aminopeptidase I activity as described above. The fractions containing dipeptidyl aminopeptidase I activity were collected, pooled, desalted and concentrated using ultrafiltration against 20 mM sodium phosphate pH 7.0 buffer.

The resultant solution was loaded onto a MonoQ 16/10 (20 ml) pre-packed column (Pharmacia Biotech AB, Uppsala, Sweden) pre-equilibrated with 20 mM sodium phosphate pH 7.0 buffer. The dipeptidyl aminopeptidase I was eluted with a 0 to 0.27 M NaCl gradient in 20 mM sodium phosphate pH 7.0 buffer. Fractions were monitored for dipeptidyl aminopeptidase

I activity as described above. The fractions between 0.200 and 0.212 M NaCl were collected, pooled, and rebuffered with 1.7 M $(\text{NH}_4)_2\text{SO}_4$ /20 mM sodium phosphate pH 7.0 buffer using ultrafiltration as described above.

The resultant solution was loaded onto a 7 x 50 mm column containing Phenyl Superose resin (Pharmacia Biotech AB, Uppsala, Sweden) pre-equilibrated with 1.7 M $(\text{NH}_4)_2\text{SO}_4$ /20 mM sodium phosphate pH 7.0 buffer solution. The dipeptidyl aminopeptidase I was eluted with a reverse 1.7 to 0 M $(\text{NH}_4)_2\text{SO}_4$ gradient in 20 mM sodium phosphate pH 7.0 buffer. Fractions were monitored for dipeptidyl aminopeptidase I activity as described above. Two fractions possessing highest activity toward Ala-Pro-*p*-nitroanilide were found to be at least 95% homogeneous based on SDS-PAGE analysis. The major band had a molecular weight of approximately 95 kDa (range of 93-96 kDa).

Example 2: Protein sequencing and amino acid analysis methods

N-terminal sequencing of the partially purified dipeptidyl aminopeptidase I described in Example 1 and a digested fragment of the dipeptidyl aminopeptidase I was performed on an Applied Biosystems 476A Protein Sequencer (Perkin Elmer/Applied Biosystems Division, Foster City, CA) with on-line HPLC and liquid phase trifluoroacetic acid (TFA) delivery. Samples of the purified dipeptidyl aminopeptidase I were transblotted onto Novex PVDF membranes (Novex, San Diego, CA) from SDS-PAGE gels and sequenced from a blott cartridge using sequencing reagents (Perkin Elmer/Applied Biosystems Division, Foster City, CA). Detection of phenylthiohydantoin-amino acids was accomplished by on-line HPLC using Buffer A containing 3.5% tetrahydrofuran in water with 15 ml of the Premix concentrate (Perkin Elmer/Applied Biosystems Division, Foster City, CA) containing acetic acid, sodium acetate, and sodium hexanesulfonate and Buffer B containing acetonitrile. Data was collected and analyzed on a Macintosh IIsi using Applied Biosystems 610 Data Analysis software.

The partially purified dipeptidyl aminopeptidase I (MonoQ peak of Example 1) was also subjected to in-gel digestion with trypsin to generate peptide fragments of the enzyme for protein sequencing. The trypsin-digested dipeptidyl aminopeptidase I was separated by SDS-PAGE electrophoresis using 8-16% Novex Tris-glycine gels (Novex, San Diego, CA). Eight Coomassie blue stained gel pieces corresponding to a molecular weight of 95 kDa were excised and washed extensively with 200 mM NH_4HCO_3 in 50% acetonitrile. The gel pieces were subsequently reduced in 1 mM dithiothreitol (DTT) at 37°C for 20 minutes and alkylated with an equal volume of 100 mM iodoacetic acid at room temperature for 20 minutes in the dark. Gel pieces were washed repeatedly with 200 mM NH_4HCO_3 in 50% acetonitrile. Supernatants were removed and gel pieces were dried on a Speed-Vac (Savant Instruments, Farmingdale, NY). The gel pieces were rehydrated in a solution containing 0.033 mg of sequencing grade modified porcine trypsin (Promega, Madison, WI) per ml of 135 mM NH_4HCO_3 . The solution was prepared by diluting 1 part of 0.1 mg of the trypsin per ml of

trypsin resuspension buffer (Promega, Madison, WI) into 2 parts of 200 mM NH_4HCO_3 . The gel pieces were incubated at 37°C for 20 hours. The peptide fragments were extracted from the gel in repetitive washes of 0.1% TFA in 60% acetonitrile for 1 hour each. The extracted peptides were dried and reconstituted in 0.05% TFA in 25% acetonitrile and then further diluted into 0.05% TFA. The peptide fragments were filtered using a Micropure 0.45 μm filter unit (Amicon, Inc., Beverly, MA). The peptide fragments were then separated by reverse-phase HPLC using a Hewlett-Packard 1090L HPLC equipped with a 2.1 x 250 mm Vydac C18-RP column (5 micron). A step gradient was used with 0.1% TFA in 80% acetonitrile as the eluant. The peptide samples were hand collected and then subjected to N-terminal sequencing.

The partially purified dipeptidyl aminopeptidase I was also subjected to cyanogen bromide to generate peptide fragments of the enzyme for sequencing. The dipeptidyl aminopeptidase I was digested with cyanogen bromide by reconstituting a dried sample of the partially purified n 70% formic acid with a few crystals of cyanogen bromide and incubating for 18 hours at room temperature in the dark. The peptide fragments were separated by SDS-PAGE electrophoresis using 10-20% Novex Tricine gels (Novex, San Diego, CA) and sequenced as described above.

N-terminal sequencing of the dipeptidyl aminopeptidase I revealed that the N-terminus was apparently blocked. A weak sequence was obtained as follows where amino acid residues in parentheses are not 100% certain and residues marked with an X could not be determined:

Peptide 1: XEGSKRLTFXETVVKQAIT(P) (SEQ ID NO:3)

The cyanogen bromide-degraded fragments had the following amino acid sequences where amino acid residues underlined matched 100% with a dipeptidyl aminopeptidase I of *Saccharomyces cerevisiae* (Anna-Arriola and Herskowitz, 1994, *Yeast* 10: 801-810; Galisson and Dujon, 1996, *Yeast* 12: 877-885):

Peptide 2: QRLPPGFSPDKKYPILFTPYGG (SEQ ID NO:4)

Peptide 3: KYIGPIK (SEQ ID NO:5)

Peptide 4: GEGSKRL (SEQ ID NO:6)

In-gel digestion with trypsin produced the following peptides where amino acid residues underlined matched 100% with the deduced amino acid sequence of the *Aspergillus oryzae* dipeptidyl aminopeptidase I nucleic acid sequence described in Example 7:

Peptide 5: XPILFTPY (SEQ ID NO:7)

Peptide 6: XVPLMPDQ(Q)GDIQYAQ (SEQ ID NO:8)

Example 3: Genomic DNA extraction

Aspergillus oryzae 1568 was grown in 25 ml of 0.5% yeast extract-2% glucose (YEG) medium for 24 hours at 37°C and 250 rpm. Mycelia were then collected by filtration through Miracloth (Calbiochem, La Jolla, CA) and washed once with 25 ml of 10 mM Tris-1 mM

EDTA (TE) buffer. Excess buffer was drained from the mycelia preparation which was subsequently frozen in liquid nitrogen. The frozen mycelia preparation was ground to a fine powder in an electric coffee grinder, and the powder was added to a disposable plastic centrifuge tube containing 20 ml of TE buffer and 5 ml of 20% w/v sodium dodecylsulfate (SDS). The mixture was gently inverted several times to ensure mixing, and extracted twice with an equal volume of phenol:chloroform:isoamyl alcohol (25:24:1 v/v/v). Sodium acetate (3 M solution) was added to the extracted sample to a final concentration of 0.3 M followed by 2.5 volumes of ice cold ethanol to precipitate the DNA. The tube was centrifuged at 15,000 x g for 30 minutes to pellet the DNA. The DNA pellet was allowed to air-dry for 30 minutes before resuspension in 0.5 ml of TE buffer. DNase-free ribonuclease A was added to the resuspended DNA pellet to a concentration of 100 µg/ml and the mixture was then incubated at 37°C for 30 minutes. Proteinase K (200 µg/ml) was added and the tube was incubated an additional one hour at 37°C. Finally, the sample was extracted twice with phenol:chloroform:isoamyl alcohol and the DNA precipitated with ethanol. The precipitated DNA was washed with 70% ethanol, dried under vacuum, resuspended in TE buffer, and stored at 4°C.

Example 4: PCR amplification of *Aspergillus oryzae* 1568 dipeptidyl aminopeptidase I

The forward degenerate oligonucleotide primer was designed to the peptide sequence DW(I/V)YEEE, a conserved motif found in most published dipeptidyl aminopeptidase I protein sequences. The reverse degenerate oligonucleotide primer was designed to a partial peptide, PPGFSDKKYP, of peptide 2 (SEQ ID NO:4) as described in Example 2. The degenerate oligonucleotide primers shown below were synthesized with an Applied Biosystems Model 394 DNA/RNA Synthesizer, according to the manufacturer's instructions, for use to PCR amplify dipeptidyl aminopeptidase I gene fragments from *Aspergillus oryzae* 1568 genomic DNA:

Forward primer: 5'-GAYTGGITITAYGARGARGAR-3' (SEQ ID NO:9)

Reverse primer: 5'-GGRTAYTTYTTRTCIGGISWRAAICCGIGG-3' (SEQ ID NO:10)

(R= A or G, Y= C or T, S= G or C, W= A or T, I= Inosine)

Amplification reactions (100 µl) were prepared using approximately 1 µg of genomic DNA isolated from an *Aspergillus oryzae* 1568 as described in Example 3 as the template. Each reaction contained the following components: 1 µg genomic DNA, 40 pmol forward primer, 40 pmol reverse primer, 200 µM each of dATP, dCTP, dGTP, and dTTP, 1 x *Taq* polymerase buffer (Perkin-Elmer Corp., Branchburg, NJ), and 2.5 Units of *Taq* polymerase (Perkin-Elmer Corp., Branchburg, NJ). The reactions were incubated in a Perkin-Elmer Model 480 Thermal Cycler programmed as follows: Cycle 1 at 94°C for 2 minutes, 45°C for one minutes, and 72°C for one minute; and cycles 2-30 each at 94°C for one minute, 45°C for

one minute, and 72°C for one minute. The reaction products were isolated on a 1% agarose gel (Eastman Kodak, Rochester, NY). An approximately 1.0 kb product band was excised from the gel and purified using GenElute spin columns (Supelco, Bellefonte, PA) according to the manufacturer's instructions. The purified PCR product was subsequently cloned into a pCRII vector (Invitrogen, San Diego, CA) and the DNA sequence was determined using lac forward and reverse primers (New England BioLabs, Beverly, MA).

A dipeptidyl aminopeptidase I gene segment consisting of approximately 321 codons (963bp) was amplified from *Aspergillus oryzae* 1568 with the dipeptidyl aminopeptidase I PCR primers described above. DNA sequence analysis shows that the amplified gene segment encoded a portion of the corresponding *Aspergillus oryzae* 1568 dipeptidyl aminopeptidase I gene. The dipeptidyl aminopeptidase I gene segment was used to probe an *Aspergillus oryzae* 1568 genomic DNA library.

Example 5: Construction of DNA libraries

A genomic DNA library was constructed in the bacteriophage cloning vector λ ZipLox (Life Technologies, Gaithersburg, MD). First, total cellular DNA was partially digested with *Tsp*509I and size-fractionated on 1% agarose gels. DNA fragments migrating in the size range 3-7 kb were excised and eluted from the gel using Prep-a-Gene reagents (BioRad Laboratories, Hercules, CA). The eluted DNA fragments were ligated with *Eco*RI-cleaved and dephosphorylated λ ZipLox vector arms (Life Technologies, Gaithersburg, MD), and the ligation mixtures were packaged using commercial packaging extracts (Stratagene, La Jolla, CA). The packaged DNA libraries were plated and amplified in *Escherichia coli* Y1090ZL cells (Life Technologies, Gaithersburg, MD). The unamplified genomic DNA library contained 3.1×10^6 pfu/ml (background titers with no DNA were 2.0×10^4 pfu/ml).

Example 6: Identification of dipeptidyl aminopeptidase I clones

Approximately 10,000 plaques from the library described in Example 5 were screened by plaque-hybridization using the dipeptidyl aminopeptidase I PCR fragment from *Aspergillus oryzae* 1568 as the probe. The DNA was cross-linked onto membranes (Hybond N+, Amersham, Arlington Heights, IL) using a UV Stratalinker (Stratagene, La Jolla, CA). The membranes were soaked for three hours at 45°C in a hybridization solution containing 5 x SSPE, 0.3% SDS, 50% formamide, and 10 μ g/ml of denatured and sheared herring sperm DNA. The dipeptidyl aminopeptidase I gene fragment isolated from the *Aspergillus oryzae* 1568 genomic DNA as described in Example 2 was radiolabeled using the Random Primed DNA Labeling Kit (Boehringer Mannheim, Mannheim, Germany), denatured by adding NaOH to a final concentration of 0.1M, and added to the hybridization solution at an activity of approximately 1×10^6 cpm per ml of hybridization solution. The mixture was incubated

overnight at 45°C in a shaking water bath. Following incubation, the membranes were washed once in 2 x SSC with 0.2% SDS at 55°C followed by two washes in 2 x SSC at the same temperature. The membranes were dried on blotting paper for 15 minutes, wrapped in SaranWrap™, and exposed to X-ray film overnight at -70°C with intensifying screens (Kodak, Rochester, NY).

Based on the production of strong hybridization signals with the probe, three plaques, designated *E. coli* DH5α MWR52A, *E. coli* DH5α MWR52B, and *E. coli* DH5α MWR52C were chosen for further study. The three plaques were purified twice in *E. coli* Y1090ZL cells and the dipeptidyl aminopeptidase I genes were subsequently excised from the λZipLox vector as pZL1-derivatives (D'Alessio et al., 1992, Focus® 14:76) using *in vivo* excision by infection of *E. coli* DH10BZL cells (Life Technologies, Gaithersburg, MD). The three plasmid containing colonies were inoculated into three ml of LB plus 50 µg/ml carbenicillin medium and grown overnight at 37°C. Miniprep DNA was prepared from each of these cultures using the Wizard 373 DNA Purification Kit (Promega, Madison, WI). The dipeptidyl aminopeptidase I encoding plasmid (pMWR52) was confirmed by DNA sequencing.

Example 7: DNA sequence analysis of *Aspergillus oryzae* 1568 dipeptidyl aminopeptidase I gene

DNA sequencing of the dipeptidyl aminopeptidase I gene contained on pMWR52 in *E. coli* DH5α MWR52 described in Example 6 was performed with an Applied Biosystems Model 377 Automated DNA Sequencer (Applied Biosystems, Inc., Foster City, CA) on both strands using the primer walking technique with dye-terminator chemistry (Giesecke *et al.*, 1992, *Journal of Virology Methods* 38: 47-60). Oligonucleotide sequencing primers were designed to complementary sequences in the dipeptidyl aminopeptidase I gene and were synthesized on an Applied Biosystems Model 394 DNA/RNA Synthesizer according to the manufacturer's instructions.

The nucleotide sequence of the gene encoding the *Aspergillus oryzae* 1568 dipeptidyl aminopeptidase I is shown in Figure 1 (SEQ ID NO:1). Sequence analysis of the cloned insert revealed an open reading frame of 2396 nucleotides (excluding the stop codon) interrupted by an 83 bp intron. The G+C content of this open reading frame was 55.3%. The deduced amino acid sequence encoded a protein of 771 amino acids (SEQ ID NO:2). Based on the rules of van Heijne (van Heijne, 1984, *Journal of Molecular Biology* 173: 243-251), the first 16 amino acids likely comprise a secretory signal peptide which directs the nascent polypeptide into the endoplasmic reticulum (boxed in Figure 1).

The amino acid sequences of the partial peptides derived from the purified dipeptidyl aminopeptidase I as described in Example 2 are underlined in Figure 1 and are consistent with

those found in the deduced amino acid sequence (SEQ ID NO:2) of the *Aspergillus oryzae* 1568 dipeptidyl aminopeptidase I cDNA.

Using the Clustal alignment program (Higgins, 1989, *CABIOS* 5: 151-153) to compare the deduced amino acid sequence of the *Aspergillus oryzae* 1568 dipeptidyl aminopeptidase I to that of a *Saccharomyces cerevisiae* dipeptidyl aminopeptidase I (Anna-Arriola and Herskowitz, 1994, *Yeast* 10: 801-810) (SEQ ID NO:11), a 23.2% identity was observed.

Example 8: Construction of a *Aspergillus oryzae* 1568 dipeptidyl aminopeptidase *Fusarium* expression vector

The coding region of *Aspergillus oryzae* dipeptidyl aminopeptidase I was amplified and the resulting fragment was cloned into pDM181 for expression in *Fusarium*. pDM181 provides the *Fusarium* trypsin (SP387) promoter and terminator, and the *bar* selectable marker gene. Specifically, the fragment was amplified by PCR using a sense primer (P1) designed to the first in-frame ATG and extending 13 bp downstream and an antisense primer (P2) designed to a region of the transcriptional stop codon and extending 10 bp downstream. To facilitate the cloning of the amplified fragment the sense and antisense primers contain a *Swa*I and a *Pac*I restriction site, respectively. The oligonucleotide primers shown below were synthesized using an ABI Model 394 DNA/RNA Synthesizer (Applied Biosystems, Inc., Foster City, CA) according to the manufacturer's instructions.

*Swa*I

P1: 5'-GATTAAATCACCATGAAGGTACGTCAATTCCACTG-3' (SEQ ID NO:12)

*Pac*I

P2: 5'-GTTAATTAATCTACTCCTCCAAGTCCTTCTTAGTCC-3' (SEQ ID NO:13)

The 50 µl PCR solution (10 mM Tris-HCl pH 8.3, 50 mM KCl, 1.5 mM MgCl₂, 0.01% w/v gelatin) contained approximately 200 ng of pMWR52 DNA, 200 µM each of dATP, dCTP, dGTP, and dTTP, and 50 pmol of each PCR primer described above. Five units of PWO polymerase (Boehringer Mannheim, Indianapolis, IN) were added and the reaction was incubated at 95°C for 3 minutes and cooled to 80°C. The reaction was then cycled 30 times, each cycle at 95°C for 30 seconds, 57°C for 1 minute, and 72°C for 1 minute, in a Perkin-Elmer 9600 Thermal Cycler. Following the last cycle, the reaction incubated for 5 minutes at 72°C.

A predicted 2.4 kb fragment was isolated by digestion with *Swa*I and *Pac*I and was cloned into pDM181 (Figure 2) digested with the same restriction endonucleases to create pMWR54 (Figure 3). To verify the fidelity of the cloned PCR fragment, the fragment was sequenced according to the method of Hattori and Sakaki (1986, *Analytical Biochemistry* 152: 232-237) using an automated Applied Biosystems Model 373A Sequencer (Applied Biosystems, Inc., Foster City, CA) according to the manufacturer's instructions.

Sequencing of the cloned dipeptidyl aminopeptidase amplified insert of pMWR54

confirmed that there were no differences in the sequence described in SEQ ID NO:1.

Example 9: Transformation of *Fusarium* CC1-3 and analysis of transformants

Fusarium strain CC1-3, a highly branched morphological mutant of *Fusarium* strain A3/5 (ATCC 20334) (Wiebe *et al.*, 1992, *Mycological Research* 96: 555-562; Wiebe *et al.*, 1991, *Mycological Research* 95: 1284-1288; Wiebe *et al.*, 1991, *Mycological Research* 96: 555-562), was grown in a liquid medium containing Vogel's salts, (Vogel, 1964, *Am. Nature* 98: 435-446), 25 mM NaNO₃, and 1.5% glucose for 4 days at 28°C and 150 rpm. Conidia were purified by filtration through 4 layers of cheesecloth and finally through one layer of Miracloth. Conidial suspensions were concentrated by centrifugation. Fifty ml of YPG medium comprised of 1% yeast extract, 2% bactopectone, and 2% glucose were inoculated with approximately 10⁸ conidia, and incubated for 14 hours at 24°C and 150 rpm. Resulting hyphae were trapped on a sterile 0.4 mm filter and washed successively with sterile distilled water and 1.0 M MgSO₄. The hyphae were resuspended in 10 ml of NOVOZYM 234™ solution (2-10 mg/ml in 1.0 M MgSO₄) and digested for 15-30 minutes at 34°C with agitation at 80 rpm. Undigested hyphal material was removed from the resulting protoplast suspension by successive filtration through 4 layers of cheesecloth and through Miracloth. Twenty ml of 1 M sorbitol were combined with the protoplast solution. After mixing, the protoplasts were pelleted by centrifugation and washed successively by resuspension and centrifugation in 20 ml of 1 M sorbitol and in 20 ml of STC (0.8 M sorbitol, 0.05 M Tris pH 8.0, 0.05 M CaCl₂). The washed protoplasts were resuspended in 4 parts STC and 1 part SPTC (0.8 M sorbitol, 40% PEG 4000, 0.05 M Tris pH 8.0, 0.05 M CaCl₂) at a concentration of 5 x 10⁷/ml.

One hundred ml of protoplast suspension were added to 10 µg of pMWR54 in polypropylene tubes (17 x 100 mm), mixed and incubated on ice for 30 minutes. One ml of SPTC was mixed gently into the protoplast suspension and incubation was continued at room temperature for 20 minutes. 12.5 ml of molten solution (cooled to 40°C) consisting of 1X Vogel's salts, 25 mM NaNO₃, 0.8 M sucrose and 1% low melting agarose (Sigma Chemical Company, St. Louis, MO) were mixed with the protoplasts and then plated onto an empty 100 mm Petri plate. Incubation was continued at room temperature for 10 to 14 days. After incubation at room temperature for 24 hours, 12.5 ml of the identical medium plus 10 mg of BASTA™ (Hoechst Schering, Rodovre, Denmark) per ml were overlayed onto the Petri plate. BASTA™ was extracted twice with phenol:chloroform:isoamyl alcohol (25:24:1), and once with chloroform:isoamyl alcohol (24:1) before use.

After two weeks, 21 transformants were apparent. A mycelial fragment from the edge of each transformant was transferred to individual wells of a 24 well plate containing Vogel's/BASTA™ medium. The medium was composed per liter of 25 g of sucrose, 25 g of Noble agar, 20 ml of 50X Vogel's salts (Vogel, 1964, *supra*), 25 mM NaNO₃, and 10 g of

BASTA™. The plate was sealed in a plastic bag to maintain moisture and incubated approximately one week at room temperature.

Example 10: Expression of *Aspergillus oryzae* 1568 dipeptidyl aminopeptidase I in *Fusarium*

A mycelial fragment from each of the 21 *Fusarium* CC1-3 transformants described in Example 9 was inoculated into 20 ml of M400Da medium composed per liter of 50 g of maltodextrin, 2.0 g of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 2.0 g of KH_2PO_4 , 4.0 g of citric acid, 8.0 g of yeast extract, 2.0 g of urea, and 0.5 ml of trace metals solution and incubated for 5 days at 30°C and 200 rpm. The medium was adjusted to pH 6.0 with 5 N NaOH. The trace metals solution was composed per liter of 14.3 g of $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, 2.5 g of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 0.5 g of $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$, 13.8 g of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 8.5 g of $\text{MnSO}_4 \cdot \text{H}_2\text{O}$, and 3.0 g of citric acid. The untransformed host was also run as a control. One ml of each culture supernatant was harvested at 5 days and stored at 4°C. Dipeptidyl aminopeptidase I activity was determined by mixing 1 µl of the enzyme supernatant with 200 µl of a Ala-Pro-pNA substrate stock solution containing 2 mg of Ala-Pro-pNA per ml of 50 mM sodium phosphate pH 7.5 and monitoring the change in absorbance at 405 nm and ambient temperature.

Culture supernatants from 19 of the 21 primary transformants of pMWR54 were positive when assayed for dipeptidyl aminopeptidase I activity.

The *Fusarium* primary transformants #1 and #16 were cultivated in 125 ml shake flasks for 5 days at 30°C in 25 ml of M400Da medium. The whole culture broths from the *Fusarium* primary transformants were filtered using a double layer of Miracloth. The filtrate was recovered and then frozen at -20°C.

Example 11: Purification of recombinant *Aspergillus oryzae* 1568 dipeptidyl aminopeptidase I produced in *Fusarium*

The recombinant dipeptidyl aminopeptidase I was purified from the *Fusarium* broth of primary transformant #1 described in Example 10. The broth (20 ml) was filtered through Nalgene filterware equipped with a 0.45 micron filter (Nalgene, Rochester, NY). The sample was diluted 10-fold using 20 mM sodium phosphate pH 7.0 buffer and concentrated using a ultrafiltration system (Amicon, Beverly, MA) utilizing a PM10 ultrafiltration membrane. The conductivity of the sample was 2.5 mS.

The sample was then loaded onto a column (XK-26) containing 60 ml of Q-Sepharose Big Beads, which had been pre-equilibrated with 400 ml of 20 mM sodium phosphate pH 7.0 buffer. The column was washed until baseline was reached. At a flow rate of 5 ml/min, the dipeptidyl aminopeptidase I was eluted with a linear gradient from 0-0.40 M NaCl in 20 mM sodium phosphate buffer pH 7.0 buffer over 10 column volumes. The dipeptidyl

aminopeptidase I eluted at ~0.24 M NaCl. SDS-PAGE was performed on the fractions active on Ala-Pro-pNA according to the following protocol. The substrate was prepared by dissolving 2 mg of Ala-Pro-pNA (Bachem, Torrance, CA) in 20 µl of DMSO. Then, 980 µl of 50 mM sodium phosphate pH 7.5 buffer was added. In a 96 well microtiter plate, 100 µl of the substrate solution was added to 100 µl of *Aspergillus oryzae* dipeptidyl aminopeptidase I (100 fold diluted in 50 mM sodium phosphate buffer pH 7.5) and the increase in absorbance at 405 nm was measured for 4 minutes using a SpectroMax 340 plate reader (Molecular Devices, Sunnyvale, Ca). The homogeneous fractions were then pooled.

Example 12: Characterization of recombinant *Aspergillus oryzae* 1568 dipeptidyl aminopeptidase I

The purified dipeptidyl aminopeptidase I described in Example 11 was characterized with respect to pH optimum, temperature stability, substrate specificity, and kinetic parameters.

The pH optimum was determined using Ala-Pro-pNA (HCl salt) as substrate in the universal buffer composed of 0.125 M citric acid, 0.125 M mono basic sodium phosphate, and 0.125 M boric acid pH was adjusted to 4.35-9.83 with 10 N NaOH, in 0.5 pH increments. The Ala-Pro-pNA substrate was prepared by dissolving 100 mg of Ala-Pro-pNA in 1 ml of DMSO and adding 20 µl of the Ala-Pro-pNA-DMSO solution to 980 µl of the universal buffer at the various pH values at ambient temperature. The assay was initiated by adding a 10 µl aliquot of the dipeptidyl aminopeptidase I diluted 20-fold in water to 200 µl of 2 mg/ml Ala-Pro-pNA at the various pH values. The absorbance at 405 nm was monitored for 5 minutes. Autohydrolysis of the substrate as a control was determined by adding 10 µl of water to 200 µl of 2 mg/ml Ala-Pro-pNA at the various pH values.

The results shown in Table 1 below demonstrated that the dipeptidyl aminopeptidase I possessed activity toward Ala-Pro-pNA as substrate over the measured pH range 4.35 to 9.83 with optimal activity at pH ~8.7. Autohydrolysis of the substrate was observed at pH values greater than 7.

Table 1

pH	Average Activity	Average Background	Avg. Activity-Background	Relative Activity
4.35	8.75 mOD/min	0 mOD/min	8.75 mOD/min	0.023
4.87	32.5	0	32.5	0.088
5.36	75.75	0	75.75	0.20
5.86	113.8	0	113.8	0.307
6.38	135.48	0	135.48	0.365
6.85	168.45	2	166.45	0.45
7.2	188.86	2	186.86	0.503
7.51	230.97	3	227.97	0.615
7.97	308.52	5.8	302.72	0.817
8.71	383.15	12.5	370.65	1
9.32	247.68	29.8	217.88	0.588
9.83	171	74	97	0.261

The temperature stability of the dipeptidyl aminopeptidase I was determined using the following protocol: 490 μ l of 50 mM sodium phosphate buffer pH 7.5 was preincubated at 37°, 45°, 55°, 60°, 65°, 70°, and 75°C for 30 minutes in a 1.7 ml Eppendorf tube. Then 10 μ l of purified dipeptidyl aminopeptidase I was added and the sample was then incubated for an additional 20 minutes. The samples were then placed on ice. Once the incubations were completed for all the temperatures, the samples were then assayed for activity using Ala-Pro-pNA as substrate.

The assay was performed by mixing 30 μ l of the incubation mixtures for the various temperatures with 200 μ l of 2 mg/ml Ala-Pro-pNA in 50 mM sodium phosphate pH 7.5 buffer) at ambient temperature. The absorbance at 405 nm was monitored for 5 minutes.

The results shown in Table 2 demonstrated that the dipeptidyl aminopeptidase I retained 90% of its activity after a 20 minute incubation at 65°C, pH 7.5.

Table 2

Temperature (°C)	Percent activity relative to 37°C
37	100
45	103
55	103
65	92.5
70	37.7
75	0.7

The relative activity of various mono-, di-, and tri-peptide para-nitroanilide substrates were assayed relative to Ala-Pro-pNA with the purified dipeptidyl aminopeptidase I diluted 100-fold in 50 mM sodium phosphate pH 7.5 buffer. Each substrate was dissolved in DMSO to a concentration of 100 mg/ml and then diluted 50 fold in 50 mM sodium phosphate pH 7.5 buffer to 2 mg/ml. The assay was performed by mixing 100 μ l of the substrate solution with 100 μ l of the dipeptidyl aminopeptidase I solution and monitoring the change in absorbance at 405 nm and ambient temperature.

The results shown in Table 3 demonstrated that the dipeptidyl aminopeptidase I preferably hydrolyzed Xaa-Pro-pNA and Xaa-Ala-pNA substrates where Xaa corresponds to any natural amino acid.

Table 3

Substrate:	Activity (mOD/min)	Percent Activity Relative to Ala-Pro-pNA
Gly-Arg-pNA	0	0
Gly-Pro-pNA	125	51.25
Arg-Pro-pNA	103	42.17
Val-Ala-pNA	49.5	20.3
Gly-Glu-pNA	<2	0
Ala-Pro-pNA	244	100
Ala-Ala-pNA	30.2	12.4
Asp-Pro-pNA	70.95	29
Leu-pNA	0	0
Ala-pNA	0	0
Ala-Ala-Pro-pNA	10.6	4.3

The kinetic parameters for various dipeptidyl aminopeptidase I substrates was determined using the following protocol. The substrates included Ala-Pro-pNA, Asp-Pro-pNA, and Ala-Ala-pNA. Purified dipeptidyl aminopeptidase I with an A_{280} of 1.521 was diluted 25-fold for Ala-Pro-pNA, 20-fold for Asp-Pro-pNA, and 10-fold for Ala-Ala-pNA in water. Each substrate was dissolved in DMSO to a concentration of 100 mg/ml and then diluted 50 fold in 50 mM sodium phosphate pH 7.5 buffer to 2 mg/ml. In a 96 well microtiter plate, 10 μ l of purified dipeptidyl aminopeptidase I was incubated with each substrate as follows except the 200 μ l substrate assay was not performed with the Asp-Pro-pNA substrate, and the absorbance at 405 nm was measured for 3 minutes:

1. 200 μ l of 2 mg/ml substrate + 0 μ l of 50 mM sodium phosphate buffer pH 7.5
2. 100 μ l of 2 mg/ml substrate + 100 μ l of 50 mM sodium phosphate buffer pH 7.5
3. 50 μ l of 2 mg/ml substrate + 150 μ l of 50 mM sodium phosphate buffer pH 7.5
4. 25 μ l of 2 mg/ml substrate + 175 μ l of 50 mM sodium phosphate buffer pH 7.5
5. 10 μ l of 2 mg/ml substrate + 190 μ l of 50 mM sodium phosphate buffer pH 7.5
6. 5 μ l of 2 mg/ml substrate + 195 μ l of 50 mM sodium phosphate buffer pH 7.5

A Lineweaver-Burke plot was constructed to determine the K_m and the k_{cat} for each substrate, using an average molecular weight of 97 kDa for the differentially glycosylated forms.

For Ala-Pro-pNA, the K_m and k_{cat} were determined to be 0.140 mM and 576.3 min^{-1} , respectively.

For Asp-Pro-pNA, the K_m and k_{cat} were determined to be 0.632 mM and 244.3 min^{-1} , respectively.

For Ala-Ala-pNA, the K_m and k_{cat} were determined to be 1.08 mM and 106.5 min^{-1} , respectively.

Example 13: Purification of *Aspergillus oryzae* 1568 aminopeptidase I

A 50 ml volume of FLAVOURZYME™ preparation described in Example 1 was centrifuged at 10,000 rpm for 10 minutes. The supernatant was filtered with a 0.2 μ m Nalgene filter and the filtrate was then diluted to 350 ml with 20 mM sodium phosphate pH 7.5 buffer. The diluted filtrate was concentrated with an Amicon ultrafiltration cell equipped with a PM10 membrane. More 20 mM phosphate pH 7.5 buffer was added and concentrated 3 more times to adjust the sample to the proper pH and conductivity.

The enzyme solution was then loaded onto a XK-26 column containing Q-Sepharose Big Beads pre-equilibrated with 20 mM phosphate buffer pH 7.5 buffer. A gradient was run from 0 to 300 mM NaCl, and then washed with 350 ml of 300 mM NaCl. Fractions were assayed for Leu-pNA activity according to the following protocol. The substrate was prepared by dissolving 2 mg of Leu-pNA (Sigma Chemical Co., St. Louis, MO) in 20 μ l of DMSO.

Then, 980 µl of 50 mM sodium phosphate pH 7.5 buffer was added. In a 96 well microtiter plate, 100 µl of the substrate solution was added to 100 µl of *Aspergillus oryzae* aminopeptidase I (100 fold diluted in 50 mM sodium phosphate pH 7.5 buffer) and the increase in absorbance at 405 nm was measured for 4 minutes using a SpectroMax 340 plate reader.

5 The most active fractions were pooled and concentrated as above using PM10 ultrafiltration.

The concentrated sample was diluted to 250 ml with 20 mM phosphate pH 7.5 buffer and loaded onto a Mono-Q 16/10 column (Pharmacia Biotech AB, Uppsala, Sweden). A gradient was run from 0 to 400 mM NaCl. Fractions were assayed for Leu-pNA activity as above. The most active and pure fractions (by SDS-PAGE) were pooled and concentrated as
10 above using using PM10 ultrafiltration.

The sample was diluted in 50 mM phosphate pH 7.0 buffer with 1.7 M ammonium sulfate and concentrated as above. This step was repeated three times. The sample was loaded onto Phenyl Sepharose column (Pharmacia Biotech AB, Uppsala, Sweden) pre-equilibrated with 50 mM sodium phosphate pH 7.0 buffer containing 1.7 M ammonium sulfate. A gradient
15 was run from 1.7 M to 0 M ammonium sulfate, and then washed with 50 mM sodium phosphate pH 7.0 buffer. Fractions were assayed for Leu-pNA activity and purity was checked by SDS-PAGE. Two small bands near 70 kDa were still present.

A 1 ml volume of the most pure fraction was concentrated with a Microcon 10 microconcentrator (Amicon, New Bedford, MA). A 100 µl volume of the concentrate was
20 loaded onto a Superose 12 column (Pharmacia Biotech AB, Uppsala, Sweden) pre-equilibrated with 20 mM phosphate buffer pH 7.0 buffer, and eluted in 30 ml of the same buffer. Only one peak was observed. SDS-PAGE analysis revealed two small bands near 70 kDa. These bands were probably an aggregation of the dipeptidyl aminopeptidase, and no further purification was necessary.

25 All active fractions from the Phenyl Sepharose column were pooled, concentrated, and desalted using PM10 ultrafiltration and 20 mM phosphate pH 7.0 buffer.

Example 14: Synergism between *Aspergillus oryzae* 1568 aminopeptidase I and *Aspergillus oryzae* 1568 dipeptidyl aminopeptidase I

30 The substrate was prepared by dissolving 2 mg of Ala-Phe-Pro-pNA (Bachem, Torrance, CA) in 20 µl of DMSO. Then, 980 µl of 50 mM sodium phosphate buffer pH 7.5 and 100 µl of ethanol were added. In a 96 well microtiter plate, 100 µl of the Ala-Phe-Pro-pNA solution was added to 100 µl of 100-fold diluted aminopeptidase I (Example 12); 100 µl of the Ala-Phe-Pro-pNA solution was added to 100 µl of 100-fold diluted dipeptidyl
35 aminopeptidase I (Example 11); and 100 µl of the Ala-Phe-Pro-pNA solution was added to a mixture of 50 µl of 100-fold diluted aminopeptidase I and 50 µl of 100-fold diluted dipeptidyl aminopeptidase I. The absorbance at 405 nm was measured for 10 minutes.

The purified aminopeptidase I was determined to have an activity of 26.822 LAPUs/ml for Leu-pNA using an extinction coefficient of 10,000 for para-nitroaniline at 405 nm. LAPU is defined as the leucine aminopeptidase activity which is determined as described in AF 298/1-GB (available on request from Novo Nordisk A/S, Denmark). The purified dipeptidyl aminopeptidase I was determined to have an activity of 11.54 DPAPU/ml for Ala-Pro-pNA, also using an extinction coefficient of 10000 for paranitroaniline. At a 100-fold dilution, aminopeptidase I exhibited less than a 2 mOD/min velocity on Ala-Pro-pNA and dipeptidyl aminopeptidase I exhibited less than a 2 mOD/min velocity on Leu-pNA. One hundred-fold diluted aminopeptidase I exhibited no activity on Ala-Phe-Pro-pNA over 10 minutes. One hundred-fold diluted dipeptidyl aminopeptidase I exhibited no activity on Ala-Phe-Pro-pNA over 10 minutes. When a mixture of 100-fold diluted dipeptidyl aminopeptidase I and 100-fold diluted aminopeptidase I were assayed with Ala-Phe-Pro-pNA, a velocity of 122 mOD/min was observed over 10 minutes.

Example 15: Preparation of protein hydrolysates with *Aspergillus oryzae* 1568 dipeptidyl aminopeptidase I

The purified dipeptidyl aminopeptidase I described in Example 11 was tested in degree of hydrolysis assays using gelatin, soy, gluten, and casein as substrates according to the following procedure.

The degree of hydrolysis (DH) assays were performed at 50°C for 18 hours as a mini-hydrolysis on a 10 ml scale using gelatin, soy bean meal tablets, wheat gluten, and sodium-caseinate at a 2% concentration adjusted to pH 7, if necessary, with no pH adjustment during hydrolysis. The hydrolyses were inactivated at 85°C for 3 minutes in a waterbath. The enzymes used were FLAVOURZYME™ with ALCALASE™ 2.4L (Novo Nordisk A/S, Bagsvrd, Denmark) and dipeptidyl aminopeptidase I with FLAVOURZYME™ and ALCALASE™ 2.4L. FLAVOURZYME™, ALCALASE™, and dipeptidyl aminopeptidase I were dosed at 3 mg, 3 mg, 0.125 mg, respectively, per 200 mg of protein per ml.

The DH, as defined by Adler-Nissen (1986, *Enzymic Hydrolysis of Food Proteins*, Elsevier Applied Science Publishers), was determined by reaction of the supernatant with OPA (ortho-phthalaldehyde, Sigma Chemical Co., St. Louis, MO). For the OPA reagent, 160 mg of OPA was dissolved in 4 ml of ethanol and transferred to a 200 ml volumetric flask containing a solution of 7.62 g of disodium tetraborate decahydrate, 200 mg of sodium dodecylsulphate, and 176 mg of dithiothreitol and the flask was filled to 200 ml with water.

A volume of 25 µl of suitably diluted supernatant was mixed with 200 µl of OPA reagent in a microtiter plate well and allowed to react for exactly 2 minutes at 25°C. The absorbance at 340 nm was measured in a microtiter plate reader and compared to the

absorbance of a 95 mM L-serine standard solution after subtraction of the blank value (water reacted with OPA-reagent). To determine the true DH, the serine equivalents measured in the supernatants were corrected with the factors suggested by *Adler-Nissen* for the trinitrobenzenesulfonic acid method (Adler-Nissen, 1979, *Agricultural and Food Chemistry* 17: 1256) which gave the same response as the described OPA method. The degree of hydrolysis was calculated on basis of the total amount of protein in the hydrolysis mixture (not on basis of soluble protein).

The results showed that dipeptidyl aminopeptidase increased the degree of hydrolysis 5% above the samples with FLAVOURZYME™ and ALCALASE™ alone for all the proteins tested.

Example 16: Increased protein solubility and release of glutamate by deamidation

Wheat gluten (WG) was obtained from Cargill (JOB 5141) and deamidated wheat gluten (DWG) was obtained from StaPro Consultancy B.V., Lemdijk 32, 9422 TH Smilde, NL. Suspensions of 8% protein were made by mixing 11 g of gluten with 89 g of water. The pH was adjusted to 6.5 with NaOH. Glutamate/aspartate specific protease (SP446), obtainable as described in WO 91/13554, or lysine/arginine specific protease (SP387) obtainable as described in WO 89/06270, was added to the suspensions. The dosage was 0.01 AU/g protein for SP446 and 0.006 AU/g protein for SP387. FLAVOURZYME™ (an non-specifically acting protease preparation available from Novo Nordisk A/S, Bagsværd, Denmark, containing endo- and exo-peptidase activities, and obtained by fermentation of *Aspergillus oryzae*) was added to some of the hydrolysates at a dosage of 20 LAPU/g protein. One LAPU (Leucine Amino Peptidase Unit) is the amount of enzyme which decomposes 1 micromole of L-leucine-p-nitroanilide per minute under the following conditions: 26 mM L-leucine-p-nitroanilide in 0.1 M Tris pH 8.0 buffer at 40°C for 10 minutes. Upon hydrolysis, p-nitroanilide is liberated turning the solution yellow which is monitored 405 nm.

The hydrolyses were carried out at 50°C without further pH adjustment for 18 hours. The enzymes were inactivated by heating at 85°C for 15 minutes. The pH was adjusted to 5 and the hydrolysates were centrifuged. The content of protein and free glutamate in the supernatant was determined.

The protein content was determined by Kjeldahl analysis, using a Kjeldahl factor of 6.25.

The content of free glutamate was determined by use of a glutamate determination kit according to the manufacturer's instructions (Boehringer-Mannheim, Indianapolis, IN). The method was adapted for use in microtiter plates.

When comparing wheat gluten (WG) to deamidated wheat gluten (DWG), the results as

shown in Table 4 demonstrated that deamidation increased the susceptibility of the gluten to specific proteases, such that more protein became soluble. By addition of FLAVOURZYME™ with a specific protease, the release of glutamate was doubled due to deamidation.

Table 4

Hydrolysate	Protein Solubility %		Glutamate Content mg/l	
	WG	DWG	WG	DWG
SP446	18	54	0	0
SP387	35	44	0	0
SP446+ FLAVOURZYME™	34	87	1000	2000

Example 17: Enzymatic deamidation and release of glutamate

Peptidoglutaminase II was produced by growing *Bacillus circulans* strain ATCC 21590 in shake flasks (400 ml) containing 200 ml of a medium composed of 1% polypeptone, 0.5% lactose, 0.025% $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.005% $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.025% KH_2PO_4 , and 17% $\text{Na}_2\text{HPO}_4 \cdot 12\text{H}_2\text{O}$ (pH adjusted to 7.2), at 30°C for 20 hours with mixing at 270 rpm. The cells were harvested by centrifugation at 4000 rpm in 1 litre flasks. The cells were then frozen.

The purification of peptidoglutaminase II from *Bacillus circulans* was performed at room temperature. The frozen *Bacillus circulans* cells were thawed and suspended in Lysis buffer (50 mM Tris/HCl; 25% (w/v) sucrose; 1 mM EDTA, pH 8.0) until a homogeneous suspension was obtained - 100 g wet cells per liter of Lysis buffer. Lysozyme (10 mg/ml) and DNase I (Sigma DN-25, 10 mg/ml) were dissolved in Lysis buffer. Then 100 ml of lysozyme solution, 10 ml of 1.0 M MgCl_2 , and 1 ml of DNase I solution were added per litre of cell suspension. The enzymes were allowed to act for 1 hour.

The suspension was filtered through a Seitz depth filter plate and the filtrate was transferred to a 10 mM $\text{KH}_2\text{PO}_4/\text{NaOH}$, pH 8.0 (Buffer A) on a Sephadex G25 column (Pharmacia). The enzyme solution was applied to a SOURCE Q column (Pharmacia) equilibrated in Buffer A and eluted with a linear NaCl gradient (0 → 500 mM) in Buffer A. Fractions from the column were analysed for Peptidoglutaminase II activity as described below and fractions with activity were pooled. The absorbance of the pooled fractions at 280 nm was 1.78, thus the protein content was estimated to 1.8 mg/ml.

The purity of the protein in the peptidoglutaminase II pool was approximately 25% as judged from a SDS-PAGE gel. Thus, the preparation contained approximately 0.5 mg/ml of

pure peptidoglutaminase II.

The peptidoglutaminase activity was determined by measuring the ammonia formed during hydrolysis of γ -carboxyamide of N-tert-Butoxycarbonyl-Gln-Pro (N-t-BOC-Gln-Pro; SIGMA No. B-4403) using the Boehringer-Mannheim kit for ammonia determination (Cat. No. 1112732). In this kit, ammonia is measured by determination of the consumption of NADH by glutamate dehydrogenase, and blanks without the addition of N-t-BOC-Gln-Pro were also applied in order to subtract the effect of other NADH consuming enzymes.

A total of 200 mg of wheat gluten protein was added to 9 ml of boiling water and after cooling, the pH was adjusted to 7.0. Then 250 μ l of the peptidoglutaminase II preparation (PEP) described above was added. The glutamate/aspartate specific protease (SP446) described in Example 16 was added in an amount of 0.04 AU/g protein, and FLAVOURZYME™ described in Example 16 was added in an amount of 20 LAPU/g protein.

Hydrolysis was allowed to proceed without pH adjustment for 18 hours at 50°C. Controls without the addition of peptidoglutaminase were also run. The hydrolysates were centrifuged and glutamate was measured as described in Example 16. The DH was determined as described in Example 15.

The results as shown below in Table 5 demonstrated that hydrolysis with the peptidoglutaminase preparation increased the DH as well as the release of glutamate.

Table 5

Hydrolysis	DH %	Glutamate mg/l
Minus PEP	40	131
Plus PEP	43	171

Deposit of Biological Materials

The following biological material has been deposited under the terms of the Budapest Treaty with the Agricultural Research Service Patent Culture Collection, Northern Regional Research Center, 1815 University Street, Peoria, Illinois, 61604, and given the following accession number:

Deposit	Accession Number	Date of Deposit
<i>E. coli</i> DH5 α pMWR52	NRRL B-21682	April 18, 1997

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

- (A) NAME: Novo Nordisk Biotech, Inc.
- (B) STREET: 1445 Drew Avenue
- (C) CITY: Davis, California
- (D) COUNTRY: United States of America
- (E) POSTAL CODE (ZIP): 95616-4880
- (F) TELEPHONE: (530) 757-8100
- (G) TELEFAX: (530) 758-0317

(ii) TITLE OF THE INVENTION: Polypeptides Having Dipeptidyl
Aminopeptidase Activity And Nucleic
Acids Encoding Same

(iii) NUMBER OF SEQUENCES: 13

(iv) CORRESPONDENCE ADDRESS:

- (A) ADDRESSEE: Novo Nordisk of North America, Inc.
- (B) STREET: 405 Lexington Avenue
- (C) CITY: New York
- (D) STATE: NY
- (E) COUNTRY: USA
- (F) ZIP: 10174

(v) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Diskette
- (B) COMPUTER: IBM Compatible
- (C) OPERATING SYSTEM: DOS
- (D) SOFTWARE: FastSEQ for Windows Version 2.0

(vi) CURRENT APPLICATION DATA:

- (A) APPLICATION NUMBER: To Be Assigned
- (B) FILING DATE: 15-May-1998
- (C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

- (A) NAME: Valeta A. Gregg
- (B) REGISTRATION NUMBER: 41,324
- (C) REFERENCE/DOCKET NUMBER: 5254.204-WO

(ix) TELECOMMUNICATION INFORMATION:

- (A) TELEPHONE: 212-867-0123
- (B) TELEFAX: 212-878-9655

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2313 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ix) FEATURE:

- (A) NAME/KEY: Coding Sequence
- (B) LOCATION: 1...2313
- (D) OTHER INFORMATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

ATG Met 1	AAG Lys	TAC Tyr	TCC Ser	AAG Lys 5	CTT Leu	CTG Leu	CTG Leu	CTC Leu	CTG Leu 10	GTC Val	AGT Ser	GTG Val	GTC Val	CAG Gln 15	GCC Ala	48
CTG Leu	GAT Asp	GTG Val	CCT Pro 20	CGG Arg	AAA Lys	CCA Pro	CAC His	GCG Ala 25	CCC Pro	ACC Thr	GGA Gly	GAA Glu	GGC Gly 30	AGT Ser	AAG Lys	96
CGT Arg	CTC Leu	ACC Thr 35	TTC Phe	AAT Asn	GAG Glu	ACC Thr	GTA Val 40	GTC Val	AAG Lys	CAA Gln	GCA Ala	ATT Ile 45	ACG Thr	CCG Pro	ACC Thr	144
TCT Ser 50	CGC Arg	TCG Ser	GTG Val	CAA Gln	TGG Trp 55	CTC Leu	TCG Ser	GGC Gly	GCA Ala	GAG Glu	GAT Asp 60	GGA Gly	TCC Ser	TAC Tyr	GTG Val	192
TAC Tyr 65	GCG Ala	GCG Ala	GAA Glu	GAC Asp 70	GGC Gly	AGT Ser	CTC Leu	ACC Thr	ATC Ile 75	GAG Glu 75	AAC Asn	ATC Ile	GTC Val	ACC Thr	AAC Asn 80	240
GAG Glu	TCA Ser	CGC Arg	ACG Thr	CTC Leu 85	ATC Ile	CCT Pro	GCG Ala	GAC Asp 90	AAG Lys	ATT Ile	CCG Pro	ACA Thr	GGG Gly	AAG Lys 95	GAA Glu	288
GCG Ala	TTC Phe	AAT Asn 100	TAC Tyr	TGG Trp	ATC Ile	CAT His	CCC Pro	GAC Asp 105	TTG Leu	TCG Ser	TCG Ser	GTG Val	CTG Leu 110	TGG Trp	GCG Ala	336
TCC Ser	AAC Asn 115	CAC His	ACC Thr	AAG Lys	CAG Gln	TAT Tyr	CGG Arg 120	CAT His	TCG Ser	TTC Phe	TTT Phe	GCC Ala 125	GAT Asp	TAT Tyr	TAC Tyr	384
GTC Val 130	CAG Gln	GAT Asp	GTG Val	GAG Glu	TCA Ser	CTC Leu 135	AAG Lys	TCC Ser	GTG Val	CCC Pro	CTG Leu 140	ATG Met	CCC Pro	GAT Asp	CAG Gln	432
GAA Glu 145	GGT Gly	GAT Asp	ATT Ile	CAA Gln	TAT Tyr	GCC Ala 150	CAA Gln	TGG Trp	AGC Ser	CCC Pro 155	GTG Val	GGC Gly	AAT Asn	ACC Thr	ATC Ile 160	480
GCT Ala	TTT Phe	GTT Val	CGC Arg	GAG Glu 165	AAT Asn	GAC Asp	CTT Leu	TAT Tyr	GTC Val 170	TGG Trp	GAT Asp	AAT Asn	GGT Gly 175	ACC Thr	GTT Val	528
ACT Thr	CGC Arg	ATT Ile	ACT Thr 180	GAT Asp	GAT Asp	GGT Gly	GGC Gly	CCC Pro 185	GAC Asp	ATG Met	TTC Phe	CAC His	GGC Gly 190	GTG Val	CCG Pro	576
GAC Asp	TGG Trp	ATC Ile 195	TAT Tyr	GAA Glu	GAG Glu	GAG Glu	ATC Ile 200	CTC Leu	GGC Gly	GAT Asp	CGC Arg	TAC Tyr 205	GCG Ala	TTG Leu	TGG Trp	624
TTC Phe 210	TCG Ser	CCA Pro	GAT Asp	GGT Gly	GAA Glu	TAT Tyr 215	CTG Leu	GCT Ala	TAC Tyr	TTG Leu	AGC Ser 220	TTC Phe	AAT Asn	GAG Glu	ACT Thr	672
GGG Gly 225	GTT Val	CCG Pro	ACC Thr	TAC Tyr	ACC Thr	GTT Val 230	CAG Gln	TAT Tyr	TAT Tyr	ATG Met 235	GAT Asp	AAC Asn	CAA Gln	GAG Glu 240	ATC Ile	720
GCT Ala	CCG Pro	GCG Ala	TAT Tyr	CCA Pro 245	TGG Trp	GAG Glu	CTG Leu	AAG Lys	ATA Ile 250	AGG Arg	TAT Tyr	CCC Pro	AAG Lys	GTG Val 255	TCG Ser	768
CAG Gln	ACG Thr	AAT Asn 260	CCG Pro	ACC Thr	GTG Val	ACG Thr	TTG Leu	AGT Ser 265	CTG Leu	CTT Leu	AAC Asn	ATC Ile	GCT Ala 270	AGC Ser	AAG Lys	816

GAG Glu	GTG Val	AAG Lys	CAG Gln	GCG Ala	CCG Pro	ATC Ile	GAC Asp	GCG Ala	TTC Phe	GAG Glu	TCA Ser	ACT Thr	GAC Asp	TTG Leu	ATC Ile	864
		275					280					285				
ATT Ile	GGC Gly	GAG Glu	GTT Val	GCT Ala	TGG Trp	CTC Leu	ACT Thr	GAT Asp	ACT Thr	CAC His	ACC Thr	ACC Thr	GTT Val	GCT Ala	GCT Ala	912
	290					295					300					
AAG Lys	GCG Ala	TTC Phe	AAC Asn	CGT Arg	GTC Val	CAG Gln	GAC Asp	CAG Gln	CAA Gln	AAG Lys	GTC Val	GTC Val	GCG Ala	GTC Val	GAT Asp	960
305					310					315					320	
ACT Thr	GCC Ala	TCG Ser	AAC Asn	AAG Lys	GCT Ala	ACT Thr	GTC Val	ATC Ile	AGC Ser	GAC Asp	CGA Arg	GAT Asp	GGG Gly	ACC Thr	GAT Asp	1008
				325					330					335		
GGA Gly	TGG Trp	CTC Leu	GAT Asp	AAC Asn	CTT Leu	CTT Leu	TCA Ser	ATG Met	AAG Lys	TAT Tyr	ATT Ile	GGC Gly	CCT Pro	ATC Ile	AAG Lys	1056
			340					345					350			
CCG Pro	TCC Ser	GAC Asp	AAG Lys	GAT Asp	GCC Ala	TAC Tyr	TAC Tyr	ATC Ile	GAC Asp	ATC Ile	TCT Ser	GAC Asp	CAT His	TCG Ser	GGA Gly	1104
		355					360					365				
TGG Trp	GCG Ala	CAT His	CTG Leu	TAT Tyr	CTC Leu	TTC Phe	CCC Pro	GTT Val	TCG Ser	GGC Gly	GGC Gly	GAA Glu	CCT Pro	ATC Ile	CCA Pro	1152
	370					375					380					
CTA Leu	ACC Thr	AAA Lys	GGC Gly	GAC Asp	TGG Trp	GAG Glu	GTC Val	ACG Thr	TCT Ser	ATT Ile	CTG Leu	AGT Ser	ATT Ile	GAT Asp	CAG Gln	1200
385					390					395					400	
GAA Glu	CGC Arg	CAG Gln	TTG Leu	GTG Val	TAC Tyr	TAC Tyr	CTG Leu	TCG Ser	ACT Thr	CAA Gln	CAC His	CAC His	AGC Ser	ACC Thr	GAG Glu	1248
				405					410					415		
CGC Arg	CAT His	CTC Leu	TAC Tyr	TCC Ser	GTC Val	TCC Ser	TAT Tyr	TCC Ser	ACG Thr	TTT Phe	GCG Ala	GTC Val	ACC Thr	CCG Pro	CTC Leu	1296
			420					425					430			
GTC Val	GAC Asp	GAC Asp	ACC Thr	GTT Val	GCC Ala	GCG Ala	TAC Tyr	TGG Trp	TCT Ser	GCT Ala	TCC Ser	TTC Phe	TCC Ser	GCG Ala	AAC Asn	1344
		435					440					445				
TCG Ser	GGC Gly	TAC Tyr	TAC Tyr	ATC Ile	CTC Leu	ACA Thr	TAC Tyr	GGA Gly	GGC Gly	CCA Pro	GAC Asp	GTA Val	CCC Pro	TAC Tyr	CAG Gln	1392
	450					455					460					
GAA Glu	CTC Leu	TAC Tyr	ACG Thr	ACC Thr	AAC Asn	AGT Ser	ACC Thr	AAA Lys	CCA Pro	CTC Leu	CGC Arg	ACA Thr	ATC Ile	ACC Thr	GAC Asp	1440
465					470				475						480	
AAC Asn	GCC Ala	AAA Lys	GTA Val	CTC Leu	GAG Glu	CAA Gln	ATC Ile	AAG Lys	GAC Asp	TAT Tyr	GCA Ala	TTG Leu	CCC Pro	AAC Asn	ATC Ile	1488
				485					490					495		
ACC Thr	TAC Tyr	TTC Phe	GAG Glu	CTT Leu	CCC Pro	CTC Leu	CCC Pro	TCC Ser	GGA Gly	GAA Glu	ACC Thr	CTC Leu	AAT Asn	GTG Val	ATG Met	1536
			500					505					510			
CAG Gln	CGC Arg	TTA Leu	CCC Pro	CCC Pro	GGG Gly	TTC Phe	TCC Ser	CCG Pro	GAT Asp	AAG Lys	AAG Lys	TAC Tyr	CCC Pro	ATA Ile	CTT Leu	1584
		515					520					525				
TTC Phe	ACC Thr	CCA Pro	TAC Tyr	GGC Gly	GGC Gly	CCA Pro	GGC Gly	GCC Ala	CAA Gln	GAA Glu	GTG Val	ACC Thr	AAG Lys	AGA Arg	TGG Trp	1632
	530					535					540					

CAA	GCC	CTG	AAT	TTC	AAG	GCC	TAT	GTC	GCC	TCC	GAC	AGC	GAA	CTC	GAG	1680
Gln	Ala	Leu	Asn	Phe	Lys	Ala	Tyr	Val	Ala	Ser	Asp	Ser	Glu	Leu	Glu	
545					550					555					560	
TAC	GTA	ACC	TGG	ACT	GTC	GAC	AAC	CGC	GGC	ACA	GGT	TTC	AAA	GGA	CGC	1728
Tyr	Val	Thr	Trp	Thr	Val	Asp	Asn	Arg	Gly	Thr	Gly	Phe	Lys	Gly	Arg	
				565					570					575		
AAG	TTC	CGC	TCC	GCC	GTC	ACG	CGC	CAA	CTC	GGC	CTC	CTC	GAA	GCA	GAA	1776
Lys	Phe	Arg	Ser	Ala	Val	Thr	Arg	Gln	Leu	Gly	Leu	Leu	Glu	Ala	Glu	
			580					585					590			
GAC	CAG	ATC	TAC	GCC	GCG	CAA	CAG	GCG	GCC	AAC	ATC	CCC	TGG	ATC	GAT	1824
Asp	Gln	Ile	Tyr	Ala	Ala	Gln	Gln	Ala	Ala	Asn	Ile	Pro	Trp	Ile	Asp	
		595					600					605				
GCA	GAC	CAC	ATC	GGC	ATC	TGG	GGC	TGG	AGT	TTC	GGA	GGC	TAC	TTG	ACC	1872
Ala	Asp	His	Ile	Gly	Ile	Trp	Gly	Trp	Ser	Phe	Gly	Gly	Tyr	Leu	Thr	
	610					615					620					
AGC	AAG	GTC	CTG	GAG	AAG	GAC	AGC	GGT	GCT	TTC	ACA	TTA	GGA	GTC	ATC	1920
Ser	Lys	Val	Leu	Glu	Lys	Asp	Ser	Gly	Ala	Phe	Thr	Leu	Gly	Val	Ile	
					630					635					640	
ACC	GCC	CCT	GTT	TCT	GAC	TGG	CGT	TTC	TAC	GAC	TCA	ATG	TAC	ACG	GAG	1968
Thr	Ala	Pro	Val	Ser	Asp	Trp	Arg	Phe	Tyr	Asp	Ser	Met	Tyr	Thr	Glu	
				645					650					655		
CGC	TAC	ATG	AAG	ACC	CTC	TCG	ACC	AAT	GAG	GAG	GGC	TAC	GAG	ACC	AGC	2016
Arg	Tyr	Met	Lys	Thr	Leu	Ser	Thr	Asn	Glu	Glu	Gly	Tyr	Glu	Thr	Ser	
			660					665					670			
GCC	GTC	CGC	AAG	ACT	GAC	GGG	TTC	AAG	AAC	GTC	GAG	GGC	GGA	TTC	TTG	2064
Ala	Val	Arg	Lys	Thr	Asp	Gly	Phe	Lys	Asn	Val	Glu	Gly	Gly	Phe	Leu	
		675					680					685				
ATC	CAG	CAC	GGA	ACG	GGC	GAC	GAT	AAC	GTC	CAT	TTC	CAG	AAC	TCG	GCT	2112
Ile	Gln	His	Gly	Thr	Gly	Asp	Asp	Asn	Val	His	Phe	Gln	Asn	Ser	Ala	
	690					695					700					
GCG	CTG	GTG	GAT	CTC	CTG	ATG	GGC	GAT	GGC	GTC	TCT	CCT	GAG	AAG	CTC	2160
Ala	Leu	Val	Asp	Leu	Leu	Met	Gly	Asp	Gly	Val	Ser	Pro	Glu	Lys	Leu	
	705				710					715					720	
CAT	TCG	CAA	TGG	TTC	ACA	GAC	TCA	GAC	CAC	GGA	ATC	AGC	TAC	CAT	GGT	2208
His	Ser	Gln	Trp	Phe	Thr	Asp	Ser	Asp	His	Gly	Ile	Ser	Tyr	His	Gly	
				725				730					735			
GGC	GGC	GTG	TTC	CTG	TAC	AAG	CAA	CTG	GCC	CGG	AAG	CTC	TAC	CAG	GAG	2256
Gly	Gly	Val	Phe	Leu	Tyr	Lys	Gln	Leu	Ala	Arg	Lys	Leu	Tyr	Gln	Glu	
			740					745					750			
AAG	AAC	CGA	CAG	ACG	CAG	GTG	CTG	ATG	CAC	CAG	TGG	ACT	AAG	AAG	GAC	2304
Lys	Asn	Arg	Gln	Thr	Gln	Val	Leu	Met	His	Gln	Trp	Thr	Lys	Lys	Asp	
		755					760					765				
TTG	GAG	GAG														2313
Leu	Glu	Glu														
		770														

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 771 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

```

Met Lys Tyr Ser Lys Leu Leu Leu Leu Leu Val Ser Val Val Gln Ala
 1      5      10      15
Leu Asp Val Pro Arg Lys Pro His Ala Pro Thr Gly Glu Gly Ser Lys
 20      25      30
Arg Leu Thr Phe Asn Glu Thr Val Val Lys Gln Ala Ile Thr Pro Thr
 35      40      45
Ser Arg Ser Val Gln Trp Leu Ser Gly Ala Glu Asp Gly Ser Tyr Val
 50      55      60
Tyr Ala Ala Glu Asp Gly Ser Leu Thr Ile Glu Asn Ile Val Thr Asn
 65      70      75      80
Glu Ser Arg Thr Leu Ile Pro Ala Asp Lys Ile Pro Thr Gly Lys Glu
 85      90      95
Ala Phe Asn Tyr Trp Ile His Pro Asp Leu Ser Ser Val Leu Trp Ala
 100      105      110
Ser Asn His Thr Lys Gln Tyr Arg His Ser Phe Phe Ala Asp Tyr Tyr
 115      120      125
Val Gln Asp Val Glu Ser Leu Lys Ser Val Pro Leu Met Pro Asp Gln
 130      135      140
Glu Gly Asp Ile Gln Tyr Ala Gln Trp Ser Pro Val Gly Asn Thr Ile
 145      150      155      160
Ala Phe Val Arg Glu Asn Asp Leu Tyr Val Trp Asp Asn Gly Thr Val
 165      170      175
Thr Arg Ile Thr Asp Asp Gly Gly Pro Asp Met Phe His Gly Val Pro
 180      185      190
Asp Trp Ile Tyr Glu Glu Glu Ile Leu Gly Asp Arg Tyr Ala Leu Trp
 195      200      205
Phe Ser Pro Asp Gly Glu Tyr Leu Ala Tyr Leu Ser Phe Asn Glu Thr
 210      215      220
Gly Val Pro Thr Tyr Thr Val Gln Tyr Tyr Met Asp Asn Gln Glu Ile
 225      230      235      240
Ala Pro Ala Tyr Pro Trp Glu Leu Lys Ile Arg Tyr Pro Lys Val Ser
 245      250      255
Gln Thr Asn Pro Thr Val Thr Leu Ser Leu Leu Asn Ile Ala Ser Lys
 260      265      270
Glu Val Lys Gln Ala Pro Ile Asp Ala Phe Glu Ser Thr Asp Leu Ile
 275      280      285
Ile Gly Glu Val Ala Trp Leu Thr Asp Thr His Thr Thr Val Ala Ala
 290      295      300
Lys Ala Phe Asn Arg Val Gln Asp Gln Gln Lys Val Val Ala Val Asp
 305      310      315      320
Thr Ala Ser Asn Lys Ala Thr Val Ile Ser Asp Arg Asp Gly Thr Asp
 325      330      335
Gly Trp Leu Asp Asn Leu Leu Ser Met Lys Tyr Ile Gly Pro Ile Lys
 340      345      350
Pro Ser Asp Lys Asp Ala Tyr Tyr Ile Asp Ile Ser Asp His Ser Gly
 355      360      365
Trp Ala His Leu Tyr Leu Phe Pro Val Ser Gly Gly Glu Pro Ile Pro
 370      375      380
Leu Thr Lys Gly Asp Trp Glu Val Thr Ser Ile Leu Ser Ile Asp Gln
 385      390      395      400
Glu Arg Gln Leu Val Tyr Tyr Leu Ser Thr Gln His His Ser Thr Glu
 405      410      415
Arg His Leu Tyr Ser Val Ser Tyr Ser Thr Phe Ala Val Thr Pro Leu
 420      425      430
Val Asp Asp Thr Val Ala Ala Tyr Trp Ser Ala Ser Phe Ser Ala Asn
 435      440      445
Ser Gly Tyr Tyr Ile Leu Thr Tyr Gly Gly Pro Asp Val Pro Tyr Gln
 450      455      460
Glu Leu Tyr Thr Thr Asn Ser Thr Lys Pro Leu Arg Thr Ile Thr Asp
 465      470      475      480

```

```

Asn Ala Lys Val Leu Glu Gln Ile Lys Asp Tyr Ala Leu Pro Asn Ile
      485      490      495
Thr Tyr Phe Glu Leu Pro Leu Pro Ser Gly Glu Thr Leu Asn Val Met
      500      505      510
Gln Arg Leu Pro Pro Gly Phe Ser Pro Asp Lys Lys Tyr Pro Ile Leu
      515      520      525
Phe Thr Pro Tyr Gly Gly Pro Gly Ala Gln Glu Val Thr Lys Arg Trp
      530      535      540
Gln Ala Leu Asn Phe Lys Ala Tyr Val Ala Ser Asp Ser Glu Leu Glu
545      550      555      560
Tyr Val Thr Trp Thr Val Asp Asn Arg Gly Thr Gly Phe Lys Gly Arg
      565      570      575
Lys Phe Arg Ser Ala Val Thr Arg Gln Leu Gly Leu Leu Glu Ala Glu
      580      585      590
Asp Gln Ile Tyr Ala Ala Gln Gln Ala Ala Asn Ile Pro Trp Ile Asp
      595      600      605
Ala Asp His Ile Gly Ile Trp Gly Trp Ser Phe Gly Gly Tyr Leu Thr
610      615      620
Ser Lys Val Leu Glu Lys Asp Ser Gly Ala Phe Thr Leu Gly Val Ile
625      630      635      640
Thr Ala Pro Val Ser Asp Trp Arg Phe Tyr Asp Ser Met Tyr Thr Glu
      645      650      655
Arg Tyr Met Lys Thr Leu Ser Thr Asn Glu Glu Gly Tyr Glu Thr Ser
      660      665      670
Ala Val Arg Lys Thr Asp Gly Phe Lys Asn Val Glu Gly Gly Phe Leu
675      680      685
Ile Gln His Gly Thr Gly Asp Asn Val His Phe Gln Asn Ser Ala
690      695      700
Ala Leu Val Asp Leu Leu Met Gly Asp Gly Val Ser Pro Glu Lys Leu
705      710      715      720
His Ser Gln Trp Phe Thr Asp Ser Asp His Gly Ile Ser Tyr His Gly
      725      730      735
Gly Gly Val Phe Leu Tyr Lys Gln Leu Ala Arg Lys Leu Tyr Gln Glu
740      745      750
Lys Asn Arg Gln Thr Gln Val Leu Met His Gln Trp Thr Lys Lys Asp
755      760      765
Leu Glu Glu
770

```

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: None

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

```

Xaa Glu Gly Ser Lys Arg Leu Thr Phe Xaa Glu Thr Val Val Lys Gln
 1           5           10           15
Ala Ile Thr Pro
      20

```

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: None

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Gln Arg Leu Pro Pro Gly Phe Ser Pro Asp Lys Lys Tyr Pro Ile Leu
 1 5 10 15
 Phe Thr Pro Tyr Gly Gly
 20

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: None

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Lys Tyr Ile Gly Pro Ile Lys
 1 5

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: None

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Gly Glu Gly Ser Lys Arg Leu
 1 5

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 8 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: None

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Xaa Pro Ile Leu Phe Thr Pro Tyr
 1 5

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 16 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: None

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Xaa Val Pro Leu Met Pro Asp Gln Gln Gly Asp Ile Gln Tyr Ala Gln
 1 5 10 15

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: None

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

```

Gly Ala Tyr Thr Gly Gly Ile Thr Ile Thr Ala Tyr Gly Ala Arg Gly
 1           5           10           15
Ala Arg Gly Ala Arg
                20

```

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: None

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

```

Gly Gly Arg Thr Ala Tyr Thr Thr Tyr Thr Thr Arg Thr Cys Ile Gly
 1           5           10           15
Gly Ile Ser Trp Arg Ala Ala Ile Cys Cys Ile Gly Gly Ile Gly Gly
                20           25           30

```

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 931 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

```

Met Ser Ala Ser Thr His Ser His Lys Arg Lys Asn Ser His Leu Phe
 1           5           10           15
Pro Gln Arg Lys Ser Ser Asn Ser Ser Met Asp Lys Pro Phe Pro
                20           25           30
Asn Asn Asp Ser Val Ala Asn Thr Asp Pro Gln Ser Asn Glu Asn Gly
                35           40           45
His Thr Ile Asn Glu Ile Arg Pro Thr Glu Ala Thr Ile Asp Val Thr
                50           55           60
Asp Val Pro Gln Thr Pro Phe Leu Gln Glu Gln Tyr Ser Met Arg Pro
        65           70           75           80
Arg Arg Glu Ser Phe Gln Phe Asn Asp Ile Glu Asn Gln His His Thr
                85           90           95
His Ser Phe Phe Ser Val Asn Lys Phe Asn Arg Arg Trp Gly Glu Trp
                100           105           110
Ser Leu Pro Glu Lys Arg Ser Tyr Val Leu Val Phe Thr Leu Ile Ala
                115           120           125
Leu Ser Val Leu Val Leu Leu Val Ile Leu Ile Pro Ser Lys Leu Leu
                130           135           140
Pro Thr Lys Ile Thr Arg Pro Lys Thr Ser Ala Gly Asp Ser Ser Leu
        145           150           155           160
Gly Lys Arg Ser Phe Ser Ile Glu Asn Val Leu Asn Gly Asp Phe Ala
                165           170           175
Ile Pro Glu Asp Thr Phe His Phe Ile Asp Pro Pro Gln Arg Leu Leu
                180           185           190
Gly Gln Asp Ser Asp Pro Gly Leu Tyr Phe Thr Thr Lys Glu Ile Asp

```

Gly	His	Thr	Asn	Phe	Ile	Ala	Lys	Gln	Leu	Phe	Asp	Glu	Thr	Phe	Glu
	210					215					220				
Val	Asn	Leu	Gly	Gly	Asn	Arg	Phe	Leu	Tyr	Glu	Gly	Val	Glu	Phe	Thr
225					230					235					240
Val	Ser	Thr	Val	Gln	Ile	Asn	Tyr	Lys	Leu	Asp	Lys	Leu	Ile	Phe	Gly
				245					250					255	
Thr	Asn	Leu	Glu	Ser	Glu	Phe	Arg	His	Ser	Ser	Lys	Gly	Phe	Tyr	Trp
			260					265					270		
Ile	Lys	Asp	Leu	Asn	Thr	Gly	Asn	Ile	Glu	Pro	Ile	Leu	Pro	Pro	Glu
		275					280					285			
Lys	Ser	Asp	Asp	Asn	Tyr	Glu	Leu	Gly	Leu	Ser	Lys	Leu	Ser	Tyr	Ala
	290					295					300				
His	Phe	Ser	Pro	Ala	Tyr	Asn	Tyr	Ile	Tyr	Phe	Val	Tyr	Glu	Asn	Asn
305					310					315					320
Leu	Phe	Leu	Gln	Gln	Val	Asn	Ser	Gly	Val	Ala	Lys	Lys	Val	Thr	Glu
				325					330					335	
Asp	Gly	Ser	Lys	Asp	Ile	Phe	Asn	Ala	Lys	Pro	Asp	Trp	Ile	Tyr	Glu
			340					345					350		
Glu	Glu	Val	Leu	Ala	Ser	Asp	Gln	Ala	Ile	Trp	Trp	Ala	Pro	Asp	Asp
		355					360					365			
Ser	Lys	Ala	Val	Phe	Ala	Arg	Phe	Asn	Asp	Thr	Ser	Val	Asp	Asp	Ile
	370					375					380				
Arg	Leu	Asn	Arg	Tyr	Thr	Asn	Met	Asn	Glu	Ala	Tyr	Leu	Ser	Asp	Thr
385					390					395					400
Lys	Ile	Lys	Tyr	Pro	Lys	Pro	Gly	Phe	Gln	Asn	Pro	Gln	Phe	Asp	Leu
				405					410					415	
Phe	Leu	Val	Asn	Leu	Gln	Asn	Gly	Ile	Ile	Tyr	Ser	Ile	Asn	Thr	Gly
			420					425					430		
Gly	Gln	Lys	Asp	Ser	Ile	Leu	Tyr	Asn	Gly	Lys	Trp	Ile	Ser	Pro	Asp
		435					440					445			
Thr	Phe	Arg	Phe	Glu	Ile	Thr	Asp	Arg	Asn	Ser	Lys	Ile	Leu	Asp	Val
	450					455					460				
Lys	Val	Tyr	Asp	Ile	Pro	Ser	Ser	Gln	Met	Leu	Thr	Val	Arg	Asn	Thr
465					470					475					480
Asn	Ser	Asn	Leu	Phe	Asn	Gly	Trp	Ile	Glu	Lys	Thr	Lys	Asp	Ile	Leu
				485					490					495	
Ser	Ile	Pro	Pro	Lys	Pro	Glu	Leu	Lys	Arg	Met	Asp	Tyr	Gly	Tyr	Ile
			500					505					510		
Asp	Ile	His	Ala	Asp	Ser	Arg	Gly	Phe	Ser	His	Leu	Phe	Tyr	Tyr	Pro
		515					520					525			
Thr	Val	Phe	Ala	Lys	Glu	Pro	Ile	Gln	Leu	Thr	Lys	Gly	Asn	Trp	Glu
	530					535					540				
Val	Thr	Gly	Asn	Gly	Ile	Val	Gly	Tyr	Glu	Tyr	Glu	Thr	Asp	Thr	Ile
545															

Gly	Gly	Lys	Gly	Trp	Ser	Phe	Arg	Ser	Trp	Ala	Arg	Glu	Lys	Leu	Gly
			740					745					750		
Tyr	Trp	Glu	Pro	Arg	Asp	Ile	Thr	Glu	Val	Thr	Lys	Lys	Phe	Ile	Gln
		755					760					765			
Arg	Asn	Ser	Gln	His	Ile	Asp	Glu	Ser	Lys	Ile	Ala	Ile	Trp	Gly	Trp
	770					775					780				
Ser	Tyr	Gly	Gly	Phe	Thr	Ser	Leu	Lys	Thr	Val	Glu	Leu	Asp	Asn	Gly
785					790					795					800
Asp	Thr	Phe	Lys	Tyr	Ala	Met	Ala	Val	Ala	Pro	Val	Thr	Asn	Trp	Thr
			805						810						815
Leu	Tyr	Asp	Ser	Val	Tyr	Thr	Glu	Arg	Tyr	Met	Asn	Gln	Pro	Ser	Glu
			820					825					830		
Asn	His	Glu	Gly	Tyr	Phe	Glu	Val	Ser	Thr	Ile	Gln	Asn	Phe	Lys	Ser
		835					840					845			
Phe	Glu	Ser	Leu	Lys	Arg	Leu	Phe	Ile	Val	His	Gly	Thr	Phe	Asp	Asp
	850					855					860				
Asn	Val	His	Ile	Gln	Asn	Thr	Phe	Arg	Leu	Val	Asp	Gln	Leu	Asn	Leu
865					870					875					880
Leu	Gly	Leu	Thr	Asn	Tyr	Asp	Met	His	Ile	Phe	Pro	Asp	Ser	Asp	His
				885					890					895	
Ser	Ile	Arg	Tyr	His	Asn	Ala	Gln	Arg	Ile	Val	Phe	Gln	Lys	Leu	Tyr
		900						905					910		
Tyr	Trp	Leu	Arg	Asp	Ala	Phe	Ala	Glu	Arg	Phe	Asp	Asn	Thr	Glu	Val
		915					920					925			
Leu	His	Leu													
		930													

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 36 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

GATTTAAATC ACCATGAAGG TACGTCAATT CCACTG

36

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 0 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GTTAATTAATCTACTCCTCCAAGTCCTTCTTAGTCC

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13 *bis*)

A. The indications made below relate to the microorganism referred to in the description on page <u>51</u> , lines <u>22-28</u>	
B. IDENTIFICATION OF Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depository institution Agricultural Research Service Patent Culture Collection (NRRL)	
Address of depository institution (<i>including postal code and country</i>) Northern Regional Research Center 1815 University Street Peoria, IL 61604, US	
Date of deposit April 18, 1997	Accession Number NRRL B-21682
C. ADDITIONAL INDICATIONS (<i>leave blank if not applicable</i>) This information is continued on an additional sheet <input type="checkbox"/>	
In respect of those designations in which a European and/or Australia Patent is sought, during the pendency of the patent application, a sample of the deposited microorganism is only to be provided to an independent expert nominated by the person requesting the sample (Rule 28(4) EPC/Regulation 3.25 of Australia Statutory Rule 1991 No. 71).	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (<i>if the indications are not for all designated States</i>)	
E. SEPARATE FURNISHING OF INDICATIONS (<i>leave blank if not applicable</i>)	
The indication listed below will be submitted to the International Bureau Later (<i>specify the general nature of the indications e.g. "Accession Number of Deposit"</i>)	

For receiving Office use only

For International Bureau use only

<input checked="" type="checkbox"/> This sheet was received with the international application	
Authorized officer	Lydell Meadows Patent Specialist IAPD-PCT Operations (800) 666-8743

<input type="checkbox"/> This sheet was received with the International Bureau on:	
Authorized officer	

Claims

What is claimed is:

- 5 1. An isolated polypeptide having dipeptidyl aminopeptidase activity, selected from the group consisting of:
 - (a) a polypeptide having an amino acid sequence which has at least 50% identity with the amino acid sequence of SEQ ID NO:2;
 - (b) a polypeptide which is encoded by a nucleic acid sequence which hybridizes under
10 medium stringency conditions with (i) the nucleic acid sequence of SEQ ID NO:1, (ii) its complementary strand, or (iii) a subsequence of SEQ ID NO:1 which encodes a polypeptide fragment which has dipeptidyl aminopeptidase activity;
 - (c) an allelic variant of (a) or (b);
 - (d) a fragment of (a), (b), or (c), wherein the fragment has dipeptidyl aminopeptidase
15 activity; and
 - (e) a polypeptide having dipeptidyl aminopeptidase activity with physicochemical properties of (i) a pH optimum in the range of from about pH 4.4 to about pH 9.8 determined after incubation for 5 minutes at ambient temperature in the presence of Ala-Pro-para-nitroanilide; (ii) a temperature stability of 90% or more, relative to initial
20 activity, at pH 7.5 determined after incubation for 20 minutes at 65°C in the absence of substrate; and (iii) an activity towards Xaa-Pro-para-nitroanilide or Xaa-Ala-para-nitroanilide wherein Xaa is selected from the group consisting of Ala, Arg, Asp, Gly, and Val.
- 25 2. The polypeptide of claim 1, comprising an amino acid sequence which has at least 50% identity with the amino acid sequence of SEQ ID NO:2.
3. The polypeptide of claim 2, comprising an amino acid sequence which has at least 60% identity with the amino acid sequence of SEQ ID NO:2.
- 30 4. The polypeptide of claim 3, comprising an amino acid sequence which has at least 70% identity with the amino acid sequence of SEQ ID NO:2.
5. The polypeptide of claim 4, comprising an amino acid sequence which has at least

80% identity with the amino acid sequence of SEQ ID NO:2.

6. The polypeptide of claim 5, comprising an amino acid sequence which has at least 90% identity with the amino acid sequence of SEQ ID NO:2.

5

7. The polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:2 or a fragment thereof.

8. The polypeptide of claim 7, comprising the amino acid sequence of SEQ ID NO:2.

10

9. The polypeptide of claim 2, which is obtained from aa *Aspergillus* strain.

10. The polypeptide of claim 9, which is obtained from an *Aspergillus oryzae* strain.

15

11. The polypeptide of claim 1, which is encoded by a nucleic acid sequence which hybridizes under medium stringency conditions with (i) the nucleic acid sequence of SEQ ID NO:1, or its complementary strand, or a subsequence thereof which encodes a polypeptide fragment which has dipeptidyl aminopeptidase activity.

20

12. The polypeptide of claim 11, which is encoded by a nucleic acid sequence which hybridizes under medium stringency conditions with (i) the nucleic acid sequence of SEQ ID NO:1 or its complementary strand.

13. The polypeptide of claim 11, which is obtained from an *Aspergillus* strain.

25

14. The polypeptide of claim 13, which is obtained from an *Aspergillus oryzae* strain.

15. The polypeptide of claim 1, which is encoded by a nucleic acid sequence which hybridizes under high stringency conditions with (i) the nucleic acid sequence of SEQ ID NO:1, or its complementary strand, or a subsequence thereof which encodes a polypeptide fragment which has dipeptidyl aminopeptidase activity.

30

16. The polypeptide of claim 15, which is encoded by a nucleic acid sequence which

hybridizes under high stringency conditions with (i) the nucleic acid sequence of SEQ ID NO:1 or its complementary strand.

17. The polypeptide of claim 15, which is obtained from an *Aspergillus* strain.

18. The polypeptide of claim 17, which is obtained from an *Aspergillus oryzae* strain.

19. The polypeptide of claim 1, which has the following physicochemical properties: (a) a pH optimum in the range of from about pH 4.4 to about pH 9.8 determined after incubation for 5 minutes at ambient temperature in the presence of Ala-Pro-para-nitroanilide; (b) a temperature stability of 90% or more, relative to initial activity, at pH 7.5 determined after incubation for 20 minutes at 65°C in the absence of substrate; and (c) an activity towards Xaa-Pro-para-nitroanilide or Xaa-Ala-para-nitroanilide wherein Xaa is selected from the group consisting of Ala, Arg, Asp, Gly, and Val.

20. The polypeptide of claim 19, which is obtained from an *Aspergillus* strain.

21. The polypeptide of claim 20, which is obtained from an *Aspergillus oryzae* strain.

22. The polypeptide of claim 1, which is encoded by the nucleic acid sequence contained in plasmid pMWR52 contained in *E. coli* NRRL B-21682.

23. The polypeptide of claim 1, which acts synergistically with an aminopeptidase to hydrolyze a polypeptide.

24. An isolated nucleic acid sequence comprising a nucleic acid sequence which encodes the polypeptide of claim 1.

25. A nucleic acid construct comprising the nucleic acid sequence of claim 24 operably linked to one or more control sequences which direct the production of the polypeptide in a suitable expression host.

26. A recombinant expression vector comprising the nucleic acid construct of claim 25, a

promoter, and transcriptional and translational stop signals.

27. A recombinant host cell comprising the nucleic acid construct of claim 25.

5 28. A method for producing the polypeptide of claim 1 comprising (a) cultivating a strain to produce a supernatant comprising the polypeptide; and (b) recovering the polypeptide.

29. A method for producing the polypeptide comprising (a) cultivating a host cell of claim 27 under conditions suitable for production of the polypeptide; and (b) recovering the
10 polypeptide.

30. A method for producing the polypeptide of claim 1 comprising (a) cultivating a homologously recombinant cell, having incorporated therein a new transcription unit comprising a regulatory sequence, an exon, and/or a splice donor site operably linked to a
15 second exon of an endogenous nucleic acid sequence encoding the polypeptide, under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.

31. A method for producing a mutant of a cell, which comprises disrupting or deleting a nucleic acid sequence encoding the polypeptide of claim 1 or a control sequence thereof,
20 which results in the mutant producing less of the polypeptide than the cell.

32. The mutant produced by the method of claim 31.

33. A method for producing a heterologous polypeptide comprising (a) culturing the
25 mutant of claim 32 under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.

34. A method for producing a hydrolysate from a proteinaceous substrate which comprises subjecting the substrate to a polypeptide of claim 1 and an endopeptidase.

5 35. The method of claim 34, wherein the hydrolysate is enriched in Ala, Arg, Asp, Gly, and/or Val.

36. A protein hydrolysate produced by the method of claim 34.

10 37. A food product comprising the protein hydrolysate of claim 36.

38. A method of obtaining from a proteinaceous substrate a hydrolysate enriched in free glutamic acid and/or peptide bound glutamic acid residues, comprising subjecting the substrate to a deamidation process and a polypeptide of claim 1.

15 39. The method of claim 38, further comprising subjecting the substrate to one or more unspecific acting endo- and/or exo-peptidase enzymes.

40. A protein hydrolysate obtained by the method of claim 39.

20 41. A food product comprising the hydrolysate of claim 40.

42. A flavor-improving composition comprising a polypeptide of claim 1 and a suitable carrier.

25 43. A pre-mix for a dough comprising a polypeptide of claim 1 and a baking ingredient.

1/4

ATGAGTCGCTTTTGTGGGCTTCGTTGCTTTCGGGCGTGTGGCTGGAGGGCGCTTGTTTCGCCGGATGAGTTCGCCGAGGATATTCAG 90
M R S L L W A S L L S G V L A G R A L V S P D E F P E D I Q
 TTGGAAGATCTGCTGGAAGGATCCCAACAGCTTGAGGACTTCGCCTATGCCTACCCCGAGCGCAATCGGTCCTTTGGTGTAAGCCAC 180
L E D L L E G S Q Q L E D F A Y A Y P E R N R V F G G K A H
 GACGACACGGTTAACTATCTCTACGAGGAGCTGAAGAAGACTGGCTACTATGATGCTCTACAAGCAGCCTCAGGTGCAGGTGTGGAGCAAT 270
 D D T V N Y L Y E E L K K T G Y Y D V Y K Q P Q V H L W S N
 GCCGACCAGACGCTCAAGGTGGGCGATGAGGAAATCGAGGGGAAGACCATGACCTACAGTCCCAGCGTCGAGGTCAACGCCGATGTAGCC 360
A D D T L K V G D E E I E A K T M T Y S P S V E V T A D V A
 GTCGTCAGAAGCTGGGATGCAGCGAGGCGGATTACCCATCCGATGTCGAGGGCAAGCTCGCCCTGATCAAGCGTGGAGAATGCCCGTTC 450
V V K N L G C S E A D Y P S D V E G K V A L I K R G E C P F
 GGCACAAAGTCGGTTCGCTGCCAAAGCCAGGCGCGGCTTCGATTGCTCTATAACAATGTGGCCGGATCCATGGCGGGCACCCCTTGGC 540
 G D K S V L A A K A K A A S I V Y N N V A G S M A G T L G
 GCGGCGCAGAGTGATAAGGACCGTATTCGGCCATTGTCGGTATCAGCTTGGAGGATGGCCAGAAGCTGATCAAGCTTGTGAGGCTGGA 630
 A A Q S D K G P Y S A I V G I S L E D G Q K L I K L A E A G
 TCGGTATCTGTGGAATCTGTGGTGGATAGTAAGCAGGAGAACCGTACGACGTATAACGTTGTCCGCAGACGAAGGGCGGCGATCCGAAC 720
 S V S V D L W V D S K Q E N R T T Y N V V A Q T K G G D P N
 AACGTCGTCGCGCTGGGTGGCCACAGGACTCAGTCGAGGGGGCCCTGGTATCAACGACGATGGCTCGGGCATTATTAGCAACTTGGTC 810
 N V V A L G G H T D S V E A G P G I N D D G S G I I S N L V

FIG.1A

2/4

ATTGCCAAAGCGCTCAGCGAGTACTCGGTCAAGAAATGCCGTGGCTTCTCTTTCTGGACAGCAGAGGAGTTCGGTCTGCTGGGCAGCAAC 900
 I A K A L T Q Y S V K N A V R F L F W T A E E F G L L G S N

 TACTACGTCTCCCATCTGAATGCCACCGAGCTGAACAAGATCCGACTGTACCTGAACCTTCGACATGATCGCCTCACCTAACTAGCCCTC 990
 Y Y V S H L N A T E L N K I R L Y L N F D M I A S P N Y A L

 ATGATCTATGACGGTGATGGATCGGCGTTCAACCAGAGCGGACCGCGGTTCCGCCAGATCGAGAAACTGTTCGAGGACTACTACGAC 1080
 M I Y D G D G S A F N Q S G P A G S A Q I E K L F E D Y Y D

 TCCATCGACCTGCCTCATATCCCCACCCAGTTTGACGGACGTTCCGACTACGAGGCGCTTATCCTGAACGGCATTCCGTCCGGTGGACTC 1170
 S I D L P H I P T Q F D G R S D Y E A F I L N G I P S G G L

 TTCACGGGCGCGAGGGCATGTCCGAAGAGAACGCAAGCCGCTGGGGAGGTCAAGCCGGGTGGCCTACGACGCCAACTACCCACGCC 1260
 F T G A E G I M S E E N A S R W G G Q A G V A Y D A N Y H A

 GCGGGAGACAACATGACCAACCTCAACCATGAAGCCTTCTGTATCAACTCCAAGCCACCGCCTTCGCCGTCCGCCACCTACGCCCAACGAC 1350
 A G D N M T N L N H E A F L I N S K A T A F A V A T Y A N D

 CTCTCTCGATCCCCAAACGGAATACCACATCCTCTTGCACCGACGAGCCCGCACCATGCGACCATTCGGCAAGAGAGCTCCGAAGACA 1440
 L S S I P K R N T T S S L H R R A R T M R P F G K R A P K T

 CACGCTCAGGTATCAGGATCCGGATGCTGGCATTCTCAAGTCGAGGCATAG 1491
 H A H V S G S G C W H S Q V E A

FIG.1B

3/4




FIG.2

4/4




Fig. 3

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 98/09629

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/57 C12N9/48 C12N1/15 C12N1/19 C12P21/06
A23J3/30

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N A23J

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	TACHI H ET AL: "AN X PROLYL DIPEPTIDYL -AMINOPEPTIDASE FROM ASPERGILLUS -ORYZAE." PHYTOCHEMISTRY (OXF) 31 (11). 1992. 3707-3709. CODEN: PYTCAS ISSN: 0031-9422, XP002047735	1-33
Y	see the whole document	34-43
A	BEAUVAIS A ET AL: "Biochemical and antigenic characterization of a new dipeptidyl- peptidase isolated from Aspergillus fumigatus." JOURNAL OF BIOLOGICAL CHEMISTRY 272 (10). 1997. 6238-6244. ISSN: 0021-9258, XP002047736 see the whole document	1-33
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

7 September 1998

Date of mailing of the international search report

21/09/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Van der Schaal, C

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/09629

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	HEYMANN E ET AL: "COMPLEMENTARY ACTION OF DIPEPTIDYL PEPTIDASE IV AND AMINOPEPTIDASE M IN THE DIGESTION OF BETA CASEIN." J DAIRY RES 53 (2). 1986. 229-236. CODEN: JDRSAN ISSN: 0022-0299, XP002052650 see page 232, last paragraph; table 2 ----	34-43
Y	CHEMICAL ABSTRACTS, vol. 124, no. 21, 20 May 1996 Columbus, Ohio, US; abstract no. 287456, TACHI, HIROSHI: "X-prolyl- dipeptidyl -aminopeptidase activity during soy sauce fermentation" XP002051874 see abstract & NIPPON JOZO KYOKAISHI (1996), 91(2), 138-40 CODEN: NJKYES; ISSN: 0914-7314, 1996, ----	34-42
Y	DATABASE WPI Section Ch, Week 9527 Derwent Publications Ltd., London, GB; Class D13, AN 95-202831 XP002047737 & JP 07 115 969 A (ASAHI KASEI KOGYO KK) , 9 May 1995 see abstract ----	34-43
Y	DATABASE WPI Section Ch, Week 9710 Derwent Publications Ltd., London, GB; Class B04, AN 97-100651 XP002076623 & CN 1 078 495 A (UNIV BEIJING) see abstract ----	34-43
Y	EP 0 480 104 A (PROTEIN TECH INT) 15 April 1992 see the whole document ----	38-41
Y	WO 96 28542 A (NOVONORDISK AS ;KAUPPINEN SAKARI (DK); SI JOAN QI (DK); SPENDLER T) 19 September 1996 see the whole document ----	42,43
P,X	BEAUVAIS A ET AL: "Dipeptidyl-peptidase IV secreted by Aspergillus fumigatus, a fungus pathogenic to humans" INFECTION AND IMMUNITY, vol. 65, no. 8, August 1997, pages 3042-3047, XP002076492 WASHINGTON US see the whole document ----	1-33
Y	see the whole document ----- -/--	34-43

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/09629

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 97 43910 A (NOVONORDISK AS ;NIELSEN PER MUNK (DK)) 27 November 1997 see claims ---	38-41
P,X	EMBL/GENBANK DATABASES, Accession no AJ002369, Sequence reference A0DPPIV 2 March 1998 DOUMAS A ET AL: "Characterization of the prolyl dipeptidyl peptidase-encoding gene (dppIV) from the koji mould Aspergillus oryzae" XP002076626	1-33
Y	see the whole document -----	34-43

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/09629

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
EP 0480104	A	15-04-1992	DK 207890 A		02-03-1991
			JP 3091445 A		17-04-1991
			NL 9001882 A		02-04-1991
			SU 1829905 A		23-07-1993
WO 9628542	A	19-09-1996	AU 4939396 A		02-10-1996
			EP 0815210 A		07-01-1998
WO 9743910	A	27-11-1997	AU 2765097 A		09-12-1997